CHEMICAL ANALYSIS OF TRITERPENE-GLYCOSIDES OF A MEDICINAL PLANT *BRYONOPSIS LACINIOSA*

N.K. SAXENA, N. SAXENA, S. SONDHIA and N. BALYANI

Department of Postgraduate Studies and Research in Chemistry, R.D. University, Jabalpur - 482001, M.P., India

Corresponding author: shobhasondia@yahoo.com

Bryonopsis laciniosa (family Cucurbitaceae) are annual herbs occurring widely in India. Ethnobotanically the whole plant is attributed with pharmacological properties. In the present studies, the bioactive molecule was isolated, purified and characterised from the whole plant. Weighed, defatted seed was subjected to methanol extraction. The isolate was purified by columns and elutes were checked for purity by thin-layer chromatography. This purified extract, called compound 1, was analysed by Infra Red Spectroscopy (IR), ¹H-Nuclear Magnetic Resonance (NMR) and ¹³C-NMR. Compound 1 was then acid hydrolysed. The glycone fraction was characterised by high performance liquid chromatography and the aglycone fraction by IR, Fast Atom Bombardment Mass Spectroscopy, ¹H-NMR and ¹³C-NMR. A tentative structure of compound 1 conforming to the data has been proposed.

EARLY DETECTION OF NEW WEEDS: HOW OFTEN TO DO SURVEILLANCE?

S.M. TIMMINS¹, S. HARRIS² and J. BROWN³

¹*Science & Research, Department of Conservation, PO Box 10-420, Wellington*

²*Harris Consulting, PO Box 70, Lyttelton*

³*Mathematics Department, Canterbury University, Christchurch*

Corresponding author: stimmins@doc.govt.nz

The Department of Conservation operates a weed surveillance system to facilitate early detection and control of new invasive weeds. Early detection minimises control costs and ecological impact. Surveillance may involve systematic checking or fortuitous finds by people engaged in other activities. Both valuable and vulnerable sites must be searched. But how often to search? Can we find new weeds early enough? When do the search costs exceed the benefits of early detection? Our model recommends how often to search for weeds in different habitats. The model considers: ability to find a new weed, rate of spread of different weeds, control costs and site biodiversity value. It suggests the surveillance interval required in forest is 1-2 years but in wetland or short vegetation it is 3 years for a short weed and 10 years for a tree. The model allows for control cost thresholds of \$500 or \$5000. In open habitat, searches must be annual to find a weed before its control cost exceeds \$500 but could be four years if a \$5000 cost threshold is acceptable. A simple cost analysis showed that it is cheaper to control environmental weeds early rather than later, i.e. weed surveillance is a good idea.