TIBOUCHINA SP. AND ANEMONE X HYBRIDA, NEW HOSTS OF STRAWBERRY LATENT RINGSPOT VIRUS IN NEW ZEALAND

J.Z. TANG, F.M. OCHOA-CORONA, B.S.M. LEBAS, D.R. ELLIOTT
and B.J.R. ALEXANDER

Ministry of Agriculture and Forestry, PO Box 2095, Auckland, New Zealand

Corresponding author: Joe.Tang@maf.govt.nz

Strawberry latent ringspot virus (SLRSV, a tentative Nepovirus), generally considered a European virus, is thought to be rare in New Zealand. In November 2003, a *tibouchina* plant (*Tibouchina* sp.) with vein chlorosis was collected from a public garden in Auckland. *Chenopodium amaranticolor* and *C. quinoa* plants inoculated with *tibouchina* leaf extracts developed systemic chlorosis with distortion and stunting of the younger leaves. *Cucumis sativum* plants exhibited interveinal chlorosis and mosaic, which matches the symptoms reported for SLRSV experimental susceptible host range. All symptomatic indicators tested positive for SLRSV by ELISA using commercially available antisera (Agdia, Inc., USA). RT-PCR using SLRSV-specific primers amplifying a 712 bp fragment of RNA2 capsid protein of SLRSV was developed. Amplified products were directly sequenced and shared 96% nucleotide identity with the published SLRSV sequence (NCBI GenBank X75165). A Japanese anemone (*Anemone x hybrida*), with leaf ringspot symptoms, collected in 1999 from an Auckland home garden had previously tested positive for SLRSV by ELISA. This too was confirmed by RT-PCR using the above primers. It is thought that this is the first report of SLRSV infecting *Tibouchina* sp. and *Anemone x hybrida* in New Zealand.

UTILISING CARROT DISKS TO SCREEN POTENTIAL BIOPESTICIDES AGAINST PRATYLENCHUS PENETRANS

L.T. DAVIS and N.L. BELL

AgResearch, Ruakura Research Centre, Private Bag 3123, Hamilton, New Zealand

Corresponding author: lee.davis@agresearch.co.nz

Pratylenchus penetrans is a migratory endoparasite of plant roots with a wide host range, including white clover, ryegrass, carrot, potato, strawberry, kiwifruit and soybean. Traditional methods for screening potential biopesticides against *P. penetrans* utilise labour intensive whole plant assays. Surface sterilised carrot disks have been used as a successful method of rearing *Pratylenchus* species, and were evaluated as a potential screening assay compared to a white clover whole plant assay. In each of the bioassays three treatments were applied: live bacteria, autoclaved bacteria and a sterile water control. Nematodes were inoculated 24 or 48 h later for the carrot (10 replicates) and white clover (12 replicates) assays respectively. The bioassays were run for eight weeks before being destructively harvested and nematodes extracted. The carrot bioassay results followed a similar trend to that of whole plants, with the bacteria treatment having the lowest nematode reproductive rate followed by autoclaved bacteria and the water control. This difference was only significant for juvenile abundance in the whole plant bioassay. A substantial labour saving favoured the carrot bioassay, largely due to reduced set up time and bioassay maintenance. The carrot bioassay is a feasible option but further development is needed to optimise the method.