DEVELOPMENT OF AN ATTRACTANT FOR THE AUSTRALIAN GUM LEAF SKELETONISER, *URABA LUGENS*, A RECENT BIOSECURITY INCURSION

A.R. GIBB¹, D.M. SUCKLING¹, S. FIELDER², J.M. ALLEN², L.E. JAMIESON³, G. CLARE³, M.L. LARSEN⁴ and G.H. WALTER⁴

¹HortResearch, PO Box 51, Lincoln, New Zealand

²HortResearch, Private Bag 11 030, Palmerston North, New Zealand

³HortResearch, Private Bag 92 169, Mt Albert, Auckland, New Zealand

⁴The University of Queensland, Brisbane, Australia

Corresponding author: agibb@hortresearch.co.nz

The Australian gum leaf skeletoniser, *Uraba lugens* Walker (Lepidoptera: Nolidae), discovered in greater Mangere, Auckland, in February 2003, is potentially a significant threat to New Zealand eucalyptus plantations and amenity plantings. Because *U. lugens* had previously been found in Auckland (2001) and Tauranga (1997 and 2000), in 2001 MAF made the decision to have its sex pheromone identified for use in detection, delimitation and eradication of any further incursions. A joint HortResearch-University of Queensland team tackled the problem. Analysis by coupled gas chromatograph-electroantennogram (GC-EAD) of the pheromone glands of *U. lugens* females revealed two major antennally active conjugated dienic compounds. Gas chromatograph-mass spectrometer analysis (GC-MS) confirmed the identity of these components. In April 2002 trapping trials were undertaken in Queensland and established that a blend of the two major components was necessary and sufficient to catch gum leaf skeletoniser males. Serendipitously, a second species of nolid moth (*Nola* sp. Leach) was also trapped in large numbers to other blends in the same trial. Deployment of monitoring traps for the latest incursion, in conjunction with further trials to refine the pheromone lure for *U. lugens*, has ensured that MAF now has an effective lure for assessing the success of any eradication programme.

CAN UNDERSTOREY DIVERSITY IN PINE PLANTATIONS AFFECT PEST LEPIDOPTERA AND THEIR NATURAL ENEMIES?

L.A. BERNDT and E.G. BROCKERHOFF

Forest Research, P.O. Box 29237, Christchurch

Corresponding author: lisa.berndt@forestresearch.co.nz

Biodiversity in plantation forests is of increasing interest to forest managers as government policy and foreign markets have increasing demands for sustainable management of these forests. Increasing indigenous biodiversity not only assists in conservation efforts, but may also play a role in ecosystem functioning, leading to a reduction in the risk of pest outbreaks. In this study we are investigating the effect of indigenous and exotic floral diversity in the understorey of a pine plantation in Canterbury on pest Lepidoptera and their parasitoids and predators. Of particular interest is the native geometrid *Pseudocoremia suavis* which has caused major defoliation in pine plantations in the past. Larval Lepidoptera were collected from pine stands with varying understorey compositions and reared to determine parasitism. Carabid beetles, which can prey on pupae on the ground, were caught using pitfall traps. The results of these experiments will be analysed and discussed.