FIELD TESTING ALTERNATIVES TO COPPER FOR CONTROLLING AVOCADO FRUIT ROTS

K.R. EVERETT, O.E. TIMUDO-TORREVILLA, G.N. HILL
and T.E. DAWSON

*The Horticulture and Food Research Institute of New Zealand Limited,
Private Bag 92169, Mt Albert, Auckland, New Zealand*

Corresponding author: Keverett@hortresearch.co.nz

ABSTRACT

Six fungicides (boscalid, boscalid/pyraclostrobin, two formulations of copper hydroxide (Kocide® Opti and Champ™ DP), dithianon and fluazinam) and one biological product (Biostart™ Target) were applied to avocado fruit in an orchard in Whangarei. Products were applied as sprays on 13 February, 23 March, 20 April, 18 June, 19 July, 21 August, 27 September and 23 October 2007. There were five replicate trees for each treatment in a completely random block design. At harvest, on 2 November, 20 fruit per tree were placed in a box, transported to Mt Albert Research Centre within 24 h, and stored at 5.5°C for 28 days. Fruit were then placed at 20°C and assessed for rots when ripe. There were too few stem-end rots for the differences between treatments to be statistically significant, but four of the fungicides significantly reduced numbers of body rots compared with the unsprayed control. These were fluazinam, boscalid/pyraclostrobin, Kocide Opti and Champ DP.

Keywords: spray trial, fungicides, rots.

INTRODUCTION

Copper use in New Zealand orchards has been of concern because of its possible detrimental effect on earthworms and microbial activity in the soil (Merrington et al. 2002; Zwieten et al. 2004) and on non-target beneficial micro-organisms and insects in the canopy (Lo & Blank 1992b; Stirling et al. 1999). Holland & Solomona (1999) surveyed the amount of copper in soil from 19 orchards in the Bay of Plenty, Gisborne, Hawke's Bay, Nelson and Otago. Of these regions, the Bay of Plenty is where avocados are grown in commercially important quantities. In the Bay of Plenty, the amount of copper in soil ranged from 80 to 210 parts per million (ppm). Severe effects of copper are expected at greater than 500 ppm copper in soil (Holland & Solomona 1999). In order to prevent the build up of copper in soils of avocado orchards to these levels, a 3-year study to find alternatives to copper fungicides was instigated. The outcomes of two strategies to achieve this aim are reported here: the first is to find alternate fungicides that are as effective as copper, the second is to find formulations of copper that are effective at lesser rates.

Last year (June 2006-July 2007) several fungicides were tested *in vitro* for effectiveness against spore germination and mycelial growth of the five fungi that most commonly cause avocado fruit rots (Everett & Timudo-Torrevilla 2007), and previously several other fungicides were similarly tested (Everett et al. 2005). The five most common fungi that cause postharvest rots on avocados are *Colletotrichum acutatum*, *C. gloeosporioides*, *Botryosphaeria parva*, *B. dothidea* and *Phomopsis* sp. (Hartill & Everett 2002; Everett et al. 2007). A fruit test was conducted to test biological products for effectiveness against these five fungi (Everett & Timudo-Torrevilla 2006).

On the basis of those results, six fungicides (boscalid, boscalid/pyraclostrobin, Champ™ DP, Kocide® Opti (previously named Kocide 3000), dithianon and fluazinam in combination with di-1-p-methene) and a biological product (Biostart™ Target) were selected for testing on an orchard.

METHODS

Fungicide application

The trial was on an avocado orchard block in Whatitiri Road, Whangarei, Northland, New Zealand. Seven products were tested, viz. boscalid, boscalid/pyraclostrobin, Champ™ DP, Kocide® Opti (previously named Kocide 3000), dithianon, fluazinam in combination with di-1-p-methene and the biological product Biostart™ Target (Table 1). Controls were unsprayed trees. Each spray treatment was applied to five replicate trees using a randomised block design. Sprays were applied at recommended rates (Table 1) at monthly intervals, a total of eight applications on 13 February, 23 March, 20 April, 18 June, 19 July, 21 August, 27 September and 23 October 2007. Products were applied using a pressurised hand gun sprayer, at approximately 8 litres/tree. Fruit were harvested on 2 November 2007, with 20 fruit being randomly selected from each of the five replicate trees, giving a total of 100 fruit/treatment.

Fruit assessment

After harvest, fruit were placed in a coolstore at the Mt Albert Research Centre of HortResearch at 5.5°C for 28 days. After that time, fruit were placed at 20°C and evaluated daily until ripe. When fruit were ripe as judged by gentle hand squeezing, they were cut in quarters, peeled and rots were assessed according to the procedures in the Avocado Industry Council assessment manual (Anon. 2003).

Statistical analysis

Results were analysed using the General Linear Model (analysis of variance) of MINITAB® (version 15.0), and means were separated using Tukey's LSD ($\alpha=0.05$). The results presented are the mean severity values = [(incidence x severity) / 100] (Fig. 1). The ORIGIN® (version 7.5) graphical package was used for drawing graphs.

TABLE 1: Application rates of products tested as spray applications for control of avocado fruit rots.

Fungicide product ¹	Active ingredient	Chemical group	Rate applied (g ai / 100 litres)
BAS 510F	boscalid	carboxamide	30 g
Pristine®	boscalid/ pyraclostrobin	carboxamide/ strobilurin	15.2/7.7 g
Champ™ DP	copper hydroxide	copper	52.5 g
Kocide® Opti	copper hydroxide	copper	27 g
Delan® 700 WG	dithianon	quinones	12.6 g
Shirlan®	fluazinam	pyridinamine	50 ml
Nufilm-17®	di-1-p-methene	terpene polymer	115.2 ml
Biostart™ Target	biological activators	ND ²	1000 ml

¹Champ is a trademark of Nufarm Americas Inc., Kocide is a trademark of Dupont, Delan and Pristine are trademarks of BASF, Shirlan is a trademark of Syngenta Group Company, Nufilm-17 is a trademark of Miller Chemical and Fertilizer Corporation, USA, and Biostart is a trademark of Biostart Limited.

²ND = not disclosed.

RESULTS

There were no statistically significant differences between treatments for stem-end rots (Fig. 1a). Four treatments significantly reduced body rots compared with levels in the untreated control (Fig. 1b). These were fluazinam, boscalid/pyraclostrobin, Champ™ DP and Kocide® Opti.

FIGURE 1: Mean severity (%) of (a) stem-end rots and (b) avocado body rots following nine field sprays at monthly intervals with test products. Fruit were harvested on 2 November 2007, coolstored for 28 days at 5.5°C and assessed after placing at 20°C when ripe. Values are means $\pm$ standard errors, and asterisks denote a significant difference from unsprayed controls (analysis of variance followed by Tukey's LSD ($\alpha=0.05$)).

The total metallic copper applied for nine applications, assuming 1500 litres/ha spray solution, was 140 g/100 litres (52.5 g ai/100 litres) = 7.1 kg/ha per season for Champ DP and 90 g/100 litres (27 g ai/100 litres) = 3.64 kg/ha per season for Kocide Opti.

DISCUSSION

In this trial, two fungicides, fluazinam and pyraclostrobin/boscalid, were as effective as copper at controlling avocado rots. A formulation of copper, Kocide Opti, was as effective as Champ DP against avocado rot fungi, at almost half the rate of elemental copper. Chemical company results (data not shown) on other crops suggest that Champ DP may also be effective at a lower rate, but this was not tested in this trial.

Thus the trial has achieved the aims of identifying alternate fungicides that were as effective as copper, and identifying copper formulations that could control avocado rots in the field using lower quantities of elemental copper.

When applied in combination with di-1-p-methene, fluazinam was an effective fungicide. This fungicide has previously shown very good efficacy in laboratory tests (Everett et al. 2005) but did not control rots when applied as an on-orchard spray. One possibility was that the formulation was damaging the skin and for this reason fluazinam was applied with di-1-p-methene as a skin protectant in this trial.

Fluazinam is a broad spectrum fungicide with a multi-site mode of action (Komyoji et al. 1995). This means that resistance is unlikely to develop in the fungal population following repeated use. There are no maximum residue limits for avocados, and residue testing now needs to be carried out before this fungicide can be registered.

Pristine is a combination fungicide of a carboxamide (boscalid) and a strobilurin (pyraclostrobin). Pyraclostrobin has a single-site mode of action, so there is a risk of resistance developing in fungal populations following repeated use. Boscalid has a different single-site mode of action and in combination with pyraclostrobin there is less chance of resistance developing than if either fungicide was applied alone (Hauke et al. 2004). When applied by itself, boscalid did not significantly reduce rots.

Although there are no maximum residue limits for avocados to pyraclostrobin or to boscalid, there is a maximum residue limit for azoxystrobin, which is also a strobilurin fungicide, for avocados in Australia. Azoxystrobin (Amistar®) has been tested on avocados in a field trial in New Zealand, and was as effective a fungicide as copper (Everett et al. 2005).

The effect of these fungicides on beneficial insects has not been investigated on avocado, but on citrus in Queensland, Australia, pyraclostrobin was reported to be less toxic to the predatory mite *Amblyseius victoriensis* Womersley (Acarina: Phytoseiidae) than the industry standard products (Miles et al. 2004). Industry standard fungicide applications included mancozeb, which has known toxicity to predators. Boscalid was non-toxic to the predacious mite *Anystis baccharum* Linnaeus (Acari: Anystidae) on apples in Canada (Laurin & Bostanian 2007). Fluazinam in combination with metalaxyl-M was slightly harmful to the beneficial arthropod *Adalia bipunctata* Linnaeus (Coleoptera: Coccinellidae), but had no effects on *Aphidius rhopalosiphii* De Stefani Perez (Hymenoptera: Braconidae) and *Episyrphus balteatus* De Geer (Diptera: Syrphidae). In contrast, copper has been observed to disrupt predation by ladybirds on citrus in New Zealand (Lo & Blank 1992a). Overseas reports are variable, and often copper is recommended as a suitable fungicide in integrated pest management programmes (Hassan et al. 1991), but it has also been recorded as having a slightly adverse effect on the beneficial insect *Cheilomenes sexmaculata* Fabricius (Coleoptera: Coccinellidae) (Krishnamoorthy et al. 2004). The effect of copper and any new fungicides on beneficial insects in avocado orchards needs to be investigated in New Zealand.

Kocide Opti has recently been registered for use on avocados in New Zealand and is available for application to orchards, to reduce the amount of elemental copper that is applied to control postharvest rots of avocados.

ACKNOWLEDGEMENTS

This study was supported by a grant from the MAF Sustainable Farming Fund, New Zealand Avocado Industry Council, BASF, Biostart, Dupont and Nufarm.

REFERENCES

- Anon. 2003. Avocado Assessment Manual, Avocado Industry Council Ltd.
- Everett KR, Boyd LM, Pak HA, Cutting JGM 2007. Calcium, fungicide sprays and canopy density influence postharvest rots of avocado. *Australasian Plant Pathology* 36(1): 22-31.
- Everett KR, Owen SG, Cutting JGM 2005. Testing efficacy of fungicides against postharvest pathogens of avocado (*Persea americana* cv. Hass). *New Zealand Plant Protection* 58: 89-95.
- Everett KR, Timudo-Torrevilla OE 2007. *In vitro* fungicide testing for control of avocado fruit rots. *New Zealand Plant Protection* 60: 99-103.
- Everett KR, Timudo-Torrevilla OE 2006. Reducing copper use in avocado orchards: results of laboratory and postharvest testing of alternate fungicides and biological products. *New Zealand Avocado Grower's Association Annual Research Report* 6: 81-90.
- Hartill WFT, Everett KR 2002. Inoculum sources and infection pathways of pathogens causing stem-end rots of 'Hass' avocado (*Persea americana*). *New Zealand Journal of Crop and Horticultural Science* 30(4): 249-260.
- Hassan SA, Bigler F, Bogenschütz H, Boller E, Brun J, Calis JNM, Chiverton P, Coremans-Pelseneer J, Duso C, Lewis GB et al. 1991. Results of the fifth joint pesticide testing programme carried out by the IOBC/WPRS-working group 'Pesticides and beneficial organisms'. *Entomophaga* 36(1): 55-67.
- Hauke K, Creemers P, Brugmans W, Laer Sv 2004. Signum, a new fungicide with interesting properties in resistance management of fungal diseases in strawberries. *Communications in Agricultural and Applied Biological Sciences* 69(4): 743-755.
- Holland P, Solomona S 1999. Copper status of orchards. *The Orchardist* 72(4): 44-45.
- Komyoji T, Sugimoto K, Mitani S, Matsuo N, Suzuki K 1995. Biological properties of a new fungicide, fluzinam. *Journal of Pesticide Science* 20(2): 129-135.
- Krishnamoorthy A, Rama N, Mani M 2004. Toxicity of pesticides to coccinellid predator *Cheilomenes sexmaculata* (Fabricius). *Insect Environment* 10(4): 183-185.
- Laurin MC, Bostanian NJ 2007. Short-term contact toxicity of seven fungicides on *Anystris baccarum*. *Phytoparasitica* 35(4): 380-385.
- Lo P, Blank R 1992a. Fungicides and insecticides disrupt predation by ladybirds on citrus. *Orchardist of New Zealand* 65(11): 14-15.
- Lo PL, Blank RH 1992b. Effect of pesticides on predation of soft wax scale by the steel-blue ladybird. *Proceedings of the 45th New Zealand Plant Protection Conference*. Pp. 99-102.
- Merrington G, Rogers SL, Zwieter Lv 2002. The potential impact of long-term copper fungicide usage on soil microbial biomass and microbial activity in an avocado orchard. *Australian Journal of Soil Research* 40(5): 749-759.
- Miles AK, Willingham SL, Cooke AW 2004. Field evaluation of strobilurins and a plant activator for the control of citrus black spot. *Australasian Plant Pathology* 33(3): 371-378.
- Stirling AM, Pegg KG, Hayward AC, Stirling GR 1999. Effect of copper fungicide on *Colletotrichum gloeosporioides* and other microorganisms on avocado leaves and fruit. *Australian Journal of Agricultural Research* 50(8): 1459-1468.
- Zwieter Lv, Rust J, Kingston T, Merrington G, Morris S 2004. Influence of copper fungicide residues on occurrence of earthworms in avocado orchard soils. *Science of the Total Environment* 329(1/3): 29-41.