THE EFFECT OF FLORAL FEEDING ON A LEAFROLLER EGG PARASITOID

P.S. STEVENS

HortResearch, Private Bag 92 169, Auckland

Trichogrammatoidea bactrae fumata is an egg parasitoid found attacking lightbrown apple moth (*Epiphyas postvittana*) and other leafrollers in New Zealand. Individual female parasitoids (< 24 hours old) were placed in a glass vial, with a newly laid *E. postvittana* egg batch, and either a clover (n=20), buttercup (n=10) or dandelion flower (n=10), or cotton wool soaked in either water (n=20) or 10% honey solution (n=10) (22°C, RH 76%, L:D 12:12). The host egg batch and food source were replaced daily for the lifetime of the parasitoid. Parasitoids provided with either water or clover had significantly lower lifetime fecundity (a mean of 2.7 and 2.6 parasitised eggs respectively) compared with parasitoids provided with honey, buttercup or dandelion flowers (14, 13.9 and 12 parasitised eggs respectively). The mean longevity of parasitoids provided with water (1.9 days) or clover flowers (1.6 days) was significantly lower than that for parasitoids provided with buttercup flowers (3.3 days). Parasitoids provided with honey or dandelion flowers had an intermediate longevity (2.7 and 2.9 days respectively) which was not significantly different from other treatments. The net reproductive rate (R_0) for females provided with water only (0.8) or clover flowers (1.8) was much lower than that of parasitoids provided with honey (6.9), buttercup flowers (9.1) or dandelion flowers (7.7).

PHENOLOGY OF THREE HOVER FLY (DIPTERA: SYRPHIDAE) SPECIES IN CANTERBURY, NEW ZEALAND

N.A. IRVIN¹, S.D. WRATTEN¹ and C.M. FRAMPTON²

¹ *Department of Entomology & Animal Ecology, P.O. Box 84, Lincoln University, NZ*

² *Centre for Computing and Biometrics, P.O. Box 84, Lincoln University, NZ*

This investigation is monitoring the populations and condition of three hoverfly species on Canterbury farmland, New Zealand. Phenology and sex ratios have been recorded from five yellow water-traps from October 15, 1993 to January 31st, 1997 and gut fullness, egg number and pollen content were recorded from September 8, 1994 to July 25, 1996. Catches of *Melanostoma fasciatum* (Macquart) and *Melangyna novaezelandiae* (Macquart) were higher during the summer months, with maximum male *M. fasciatum* catches at 0.7 per trap/day in January, 1997. Catches of female *Eristalis tenax* L. peaked at 0.24 in August, 1994. Gut fullness and egg number in *M. novaezelandiae* and *M. fasciatum* were low from June to September, 1995, a pattern opposite to that of *E. tenax*. *M. novaezelandiae* had a higher average gut fullness than did *M. fasciatum* and for all species combined, gut fullness was significantly higher in gravid females compared with non-gravid females, and males. Thirty-nine different pollen types were recorded over all. Male *E. tenax* consumed the widest range of pollen in summer and autumn whereas females consumed the widest range in winter and spring, a pattern opposite to that of *M. novaezelandiae*. The value of the work in possible population enhancement programmes for hover flies in crops is discussed.