CHEMICAL CONTROL OF TREE LUPIN

By: C. Christensen.
Foxton.

* * *

Today more than ever our national economy and moral obligations overseas demand that we utilise all available land in the production of food and clothing. Consequently any infestation of fertile grazing lands by weeds must be viewed seriously in the light of future and present production. It was this factor, that ultimately led me to the use of hormone sprays for the control and eradication of lupin.

I am farming a property of 133 acres at Oroua Downs near Foxton which was badly over-run with lupins. The land is mostly flat with a good covering of black soil while the few ridges are stable and provide quite a fair growth in those times when rainfall is sufficient.

My first approach to the problem of controlling lupin was by crushing with a crusher bar or with a Cambridge roller. I found this method fairly efficient for "Old Man" lupin but of course this still left the regrowth and seedlings to be dealt with later. I found this still to be the case even after ploughing and sowing in Chou Moellier followed by Black Barley and then grass. The seedlings quickly gained control of the pasture until practically no grazing was left. On these paddocks I used the mower with varying success but was far from satisfied with the results. Pulling and grubbing were also tried but proved too big a job and too expensive if hired labour was obtained. It was becoming increasingly apparent at this stage that I was not making any marked progress towards complete eradication. It was this fact that led me to investigate the comparatively new method of control by means of hormone sprays.

As a result in April 1949 I had a trial area laid down, using a 2,4-D concentrate at the rate of 1 quart per acre. This was a demonstration and was done with hand wands. The complete kill obtained encouraged me to purchase my own equipment and I commenced my spraying campaign at Christmas 1949. The class of lupin first treated was mature on fence lines and seedlings and regrowth in pastures. As was the case in the trial area 2,4-D concentrate was used at the same rate; the only difference being that the spraying was done through size 46 nozzles at a pressure of 50 lbs. P.S.I. on pastures and 35 lbs. P.S.I. on fence lines. As was the case in the trial area the results were again very pleasing. The procedure I now follow is to crush in late winter and spray on the regrowth and seedlings in the following January. This year I had the best kill I have experienced and better than I have seen elsewhere. Of course I have had the usual teething troubles and have not always had such excellent results but I am convinced that as my efficiency and technique improve so will the degree of the kill. Perhaps it would be as well to describe here just what my equipment consists of and the methods I have found most successful in actually applying the spray. In the first case an angle iron and hardwood frame was made to carry a 44 gallon drum across the back of the tractor. The boom we made up has 7 nozzles at 14 inch intervals. It is attached behind the container and can be raised or lowered. The pressure to nozzles is supplied by means of a pump specially made for the purpose. In the first place it was driven direct from the power take off, but I found that sufficient pressure at the nozzles could only be obtained at high engine revs. To overcome this I fitted a pulley to P.T.O. and drove the pump with a V-belt. On the front of the tractor I still have the bumper bar which is used when crushing. This I consider important as it helps the complete coverage of the plants by lowering all lupin to an almost even level. During the actual spraying a slow steady speed should be maintained - not more than three miles per hour. A near calm day is best for an even coverage and I have found it better to travel across any wind or drift starting from the windward side. The proportion of 2,4-D and water I now use has been found after many trials. I have tried more and less of each but think a quart of chemical and 50 gallons of water at 50 lbs. P.S.I. pressure gives the best results. This works out at 25/- per acre for chemicals plus the usual running expenses of the tractor.

The economics of spraying with a hormone type spray must be given serious consideration as, in general, the type of land infested with lupins

is of a poorer type.

A few of what I consider the more important points for successful spraying can be summarised as follows:-

- (1) Use 1 quart of concentrate and 50 gallons of water.
- (2) The bumper bar in front of the tractor should be used. The one I use is the same width as the boom.
- (3) Pressure should be maintained constantly at 50 lbs. P.S.I.
- (4) Land speed should be regular and not exceed 3 miles per hour.
- (5) Travel so that drift is onto the unsprayed portion.
- (6) Pasture should be closed after spraying to allow a good coverage of grass to take place.

In conclusion I feel convinced that there is a great future in the use of hormone sprays not only for the control of lupin but for all weed spraying in general providing always that the operation is planned to ensure careful application and costs watched.

* * *