EXPERIMENTAL WORK WITH PHENOXY-BUTYRICS

By F. A. MEEKLAH, Invermay Research Station, Mosgiel

SUMMARY

A SERIES of trials was conducted involving the use of gamma phenoxy-butyric acids. The amine salt of dinoseb was the most efficient treatment applied to a crop of peas intended for canning. MCPB gave significant yield increases, but 2lb. acid equivalent per acre caused slight taint. 2,4-DB and MCPB gave equal control of Californian thistle with MCPA the most efficient. The sodium salt of 2,4-DB, the butyl ester of 2,4-DB, and the ethyl ester of 2,4-D gave better control of broad-leaved docks than the sodium salt of MCPB and the butyl ester of MCPB and the potassium salt of MCPA. The sodium salt of 2,4-DB had no significant effect on seedling lucerne. The incidence of frost increased hormone damage. The sodium salt of MCPB and the butyl ester of MCPB caused no major changes in a 2-month-old sward. The sodium salt of MCPB applied at three stages to established pasture resulted in only minor reductions in yield compared with the ethyl ester of 2,4-D and the potassium salt of MCPA. Except for the summer application the potassium salt of MCPA was more phytotoxic to white clover than the ethyl ester of 2,4-D.

INTRODUCTION

Since Professor Wain's first announcement of his discoveries in 1954 with the gamma phenoxy-butyrics, we have come a long way so far as the research and development of these compounds are concerned. Nevertheless, there still remains much to be investigated and this paper will deal with a group of field trials incorporating gamma-butyrics, which were carried out at Invermay Research Station to assist in their evaluation under South Island conditions. It is important to note with the whole of this group that growing conditions throughout the season were exceptionally good. All dosages referred to in this paper are in lb. acid equivalent per acre.

A. CONTROL OF WEEDS IN PEAS

The sodium salt of MCPB at 1 and 2lb./acre was compared with 1.33lb. of the triethanolamine salt of dinoseb on a crop of Victory Freezer peas grown for processing. All treatments were applied in 33 gallons of water/acre and were replicated 6 times in a randomised block with a plot size of 20ft. x 7ft., containing 6 x 14in. rows per plot. The peas were 4 to 6in. high with 3 to 4 nodes per plant. Weather conditions, cloud five-eighths, air temperature 75 degrees F., soil temperature 67 degrees F. at 3in.

The weed population was mainly *Polygonum* spp with some fat-hen (*Chenopodium album*). Weed infestation was relatively even throughout the trial. Visual gradings of weed control were as follows:—

Dinoseb: 95 per cent. control. Some slight scorch occurred to lower leaves of peas.

MCPB 1lb. and 2lb.: There was little difference between the two rates, both giving 46 per cent. control of weeds. Peas appeared to be slightly depressed initially with some slight epinasty of the leaf stems, giving a typical droopy appearance, but at a month no difference from control could be detected.

Yields (see Table 1): All treatments gave a significantly higher yield than control. The dinoseb plots showed a 50 per cent. increase in yield, were the easiest to harvest, and ripened 2 days before other treatments. MCPB at both rates increased yields by 20 per cent., made harvesting slightly easier than control, and ripened at the same time as control.

Palatability: In tasting tests on blanched and unblanched samples only MCPB at 2lb. caused a slightly undesirable flavour in the case of the blanched sample (See Table 2).

B. CONTROL OF CALIFORNIAN THISTLE (*CIRSIIUM ARVENSE*)

The potassium salt of MCPA, the sodium salt of MCPB, and the sodium salt of 2,4-DB all at 1lb./acre were compared on thistles infesting a second-year pasture. All treatments were applied in 10 gallons of water/acre, in a randomised block of three replicates. Weather conditions at spraying were good with an air temperature of 76 degrees F. and soil temperature of 64 degrees F.

Results: Thistle control was estimated by visual grading. MCPA gave the best control; 90 per cent. 3 months after spraying. It was the quickest acting and the effects were more severe at all stages.

MCPB was initially the slowest-acting chemical in the trial. It gave 70 per cent. control at 3 months, though thistle control appeared somewhat variable due to some affected plants still being alive.

2,4-DB also showed 70 per cent. control at 3 months. The effects were also variable, some affected plants being still alive at this stage, though flowering was prevented on all plants.

It should be stressed that this is an interim report, the value of any particular herbicide being shown by the presence or absence of regrowths next spring.

C. CONTROL OF BROAD-LEAVED DOCK (*RUMEX OBTUSIFOLIUS*)

An area of waste ground on which the major plant population was mature broad-leaved docks was treated with six herbicides, the potassium salt of MCPA, the ethyl ester of 2,4-D, the sodium salt of MCPB, the butyl ester of MCPB, the sodium salt of 2,4-DB, and the butyl ester of 2,4-DB. All treatments were replicated four times in a randomised block, and all were applied at 1lb. acid equivalent per acre in 15 gallons of water. Regrowths were sprayed again in the autumn using the same formulations.

Results: Dock control was estimated by visual grading (see Table 3). Variation between replicates of any particular treatment was small, with the exception of MCPA. Poorest control was almost consistently shown by the sodium salt of MCPB, with some slight improvement by the butyl ester formulation. MCPA gave good initial control, but regrowth was extensive. The ethyl ester of 2,4-D was initially good, but regrowth was strong. The sodium salt of 2,4-DB was generally more efficient, but the amount of regrowth showing at 3 months was still high. A small increase in efficiency was shown by the butyl ester, giving good initial knockdown and small subsequent regrowth. This must be considered as a progress report, the extent of regrowth next spring providing the most important part of the picture.

D. TOLERANCE OF SEEDLING LUCERNE

The sodium salt of 2,4-DB at 1lb. and 2lb./acre was compared with 1.33lb. (phenol equivalent) of the amine salt of dinoseb when applied to N.Z. certified lucerne 1½ in. to 2 in. high, at the 2- to 3-trifoliate leaf stage. Plot sizes were 5ft. x 30ft. with the treatments replicated eight times in a randomised block layout. All treatments were applied in 23

gallons of water/acre; the weather at spraying was humid and overcast, air temperature 70 degrees F., and soil temperature 70 degrees F. at 3in. A shower of rain occurred an hour after spraying, with further showers 8 hours later. Commencing 2 days after spraying all plots were hand weeded and all weeds, affected or otherwise, were removed. Weed control assessments made before the hand weeding of the final blocks 11 days after spraying were as follows:—

2,4-DB 1lb. and 2lb. showed no difference between rates. Both gave excellent control of fat-hen, redshank (*Polygonum persicaria*), and seedling docks (*Rumex obtusifolius*). There was a fair control (mainly stunting) of cornbind (*Polygonum convolvulus*), wireweed (*P. aviculare*), and chickweed (*Stellaria media*) but poor control of shepherd's purse (*Capsella bursa-pastoris*) though some leaf roll occurred.

Dinoseb (amine): 100 per cent. control of weeds.

Effect of treatment on dry matter production (Table 4): Two yield cuts were taken and differences between treatments were small. The 1lb. and 2lb. rates of 2,4-DB gave a 7 per cent. reduction at the first cut, and dinoseb a 5 per cent. reduction. Differences were even smaller at the second cut, and at neither cut were the differences significant. Leaf abnormalities due to the 2,4-DB were only slight, consisting of the involute rolling of one or two sets of leaves typical of gamma butyric response.

Varietal susceptibility to 2,4-DB: 1lb. of the sodium salt of 2,4-DB was applied to a February-sown block of 10 lucerne varieties on 11/4/57. Stages of growth varied, but the more erect varieties were 2in. to 3in. high. Unfortunately ground frost was recorded on 21 and 22 March 1957, 31 March, and 7 April. The two early frosts caused some degree of leaf damage, the severity varying according to variety. This damage was estimated by visual grading on 8 April. Varietal susceptibility to 2,4-DB was noted 3 weeks after spraying and was graded according to damage on 3 May (see Table 5). It was found that there was a significant correlation between frost and hormone damage, and it may be postulated that frost predisposes the plant to injury by hormone weedkillers. An interesting point arising from this observation is that working with cereals, Anderson in Denmark (1) does not appear to agree with Rademacher in Germany (2) on the theory that the incidence of frost increased damage by 2,4-D; possibly Anderson was working with a frost-resistant variety or selection. Though the combination of frost and hormone damage appeared severe on a variety such as Hunter River, the hormone damage did not amount to more than the involute rolling of those sets of leaves which would be in the primordial stage at the time of spraying. The whole series of assessments for hormone damage were based on the frequency and degree of this leaf rolling. Symptoms such as a crinkling of the leaf and various degrees of chlorosis were noted at the assessment for frost and thus were not taken into account.

E. TOLERANCE OF YOUNG PASTURE

A block of November-sown pasture was sprayed when approximately 6in. high, with white clover in the 4 to 5 trifoliate leaf stage, and red clover showing 6 to 7 trifoliate leaves and just starting to tiller. Spraying was delayed until this stage by adverse weather conditions and subsequently conditions were such that the initial yield cut could not be obtained and the trial was cross trimmed. Yields were obtained at the two subsequent

cuts. Two formulations were used in this trial, the sodium salt and the butyl ester of MCPB, both at 1lb. and 2lb./acre, replicated five times and applied in 19 gallons of water/acre. Conditions at spraying were hot with an air temperature of 73 degrees F. and a soil temperature of 63 degrees F. at 3in.

Results: Differences due to treatment in this trial were so small that definite trends were difficult to detect with normal pasture evaluation techniques. Yield differences at the first cut (a month after spraying) were relatively small and were not statistically significant. Point analysis of the two high rates and controls was carried out 6 weeks after spraying, at 500 points per treatment, and showed little difference between treatments. There was no difference in clover content expressed either as actual hits, percentage sward, or percentage cover. Actual hits showed less grass in the butyl ester plots. Percentage sward and percentage cover showed a slight increase in grass in the sodium salt plots.

A second yield cut taken 12 weeks after spraying again revealed small differences, the largest being a 7 per cent. reduction in yield with 2lb. of the butyl ester, and again differences were not significant. Herbage dissection of the high rates and controls at both yield cuts (Table 6) suggested that the sodium salt was more severe on red clover than the butyl ester. This resulted initially in a slightly quicker establishment of white clover at the first cut, and a distinct gain in grass production at the second cut, while the butyl ester plots had lower grass production due to the increased competition from red clover. It could be possible that the ester is more phytotoxic to the grass than the sodium salt. The apparent lesser phytotoxicity to clovers by the butyl ester is unexpected when one remembers that the use of ester formulations of the corresponding phenoxy-acetic acids increases clover damage. One possible explanation is that more scorch on the red clover was noted in the butyl ester plots, and it is suspected that this quick destruction of small areas of leaf tissue has led to a decreased uptake of the herbicide, resulting ultimately in less over-all phytotoxicity.

F. CLOVER TOLERANCE TRIAL

To compare the tolerance of clovers in an established pasture to the sodium salt of MCPB, the potassium salt of MCPA and the ethyl ester of 2,4-D applied at different times of the year a trial was laid down on an established pasture in which white clover amounted to approximately 50 per cent. of the sward, grass 49 per cent., red clover 1 per cent. and no weeds. The two phenoxy-acetic acids were applied at 1lb., with MCPB at 2lb./acre, all in approximately 20 gallons of water/acre. Treatments were replicated four times in a randomised block. All were applied to clovers about 3½in. high at the flush of spring growth, the peak of summer growth, and at the flush of autumn clover growth.

Results: The initial reaction of the clovers to the phenoxy-acetics was very marked at all seasons, resulting in severe epinasty to leaf stems, and some scorch and subsequent death of numerous leaves. Conversely, it was difficult at all times to detect the effects of MCPB.

Spring application: Two weeks after spraying MCPA had reduced dry matter production by 22 per cent., 2,4-D 16 per cent., and MCPB 9 per cent. No differences were evident 2 months after spraying. At 3 months MCPA had improved above control, but 2,4-D was not consistent. At the end of the season there were only minor differences in total production.

Losses in production were due to clover suppression. Herbage dissection and field observation showed 2,4-D to be the most severe on white clover, but point analysis did not confirm any differences between MCPA and 2,4-D. Red Clover was eliminated from the sward by 2,4-D.

Little difference from controls could be detected with MCPB. Herbage dissection throughout the season showed that the slowest recovery from clover suppression occurred with MCPA, being 20 per cent. less in clover content than controls 2 months after spraying, and 10 per cent. less at 4 and 5 months after spraying. Clover recovery in the 2,4-D plots was somewhat faster, only a 10 per cent. suppression showing 2 months after treatment, with a slight increase (3 per cent.) over controls at 5 months.

Summer application: At the first cut following spraying, 2,4-D caused yield reductions as great as MCPA, 19 per cent. and 18 per cent. respectively, with MCPB causing virtually no reduction. Again the loss in production was due to the suppression of the clover fraction. Herbage dissection and field observation showed a similar reduction by the phenoxy-acetics of white clover by about 14 per cent. Point analysis 2 months later showed that only 2,4-D still gave any appreciable reduction in white clover content and complete elimination of red clover. MCPA showed a slight reduction in white clover and a slight increase in red clover. MCPB gave much the same picture except for a slight reduction in grass content. Field observation at this time could detect no difference between MCPA and 2,4-D nor between MCPB and control; and no differences at all at 3 months.

Dry matter production was still depressed by 2,4-D at 2 months, but at 3½ months both 2,4-D and to less degree MCPA were producing more than control.

Autumn application: The reaction of the sward to herbicides applied at this time was most severe. At 7 weeks after spraying MCPA had reduced yields by 42 per cent. with 2,4-D 28 per cent.; MCPB caused only 6 per cent. reduction. Field observations showed a more severe suppression of clover by MCP than 2,4-D. Herbage dissection showed that MCPA reduced clovers by 30 per cent., 2,4-D by 20 per cent., with MCPB showing little change with only 5 per cent. reduction. The results of point analysis were not available at the time of preparing this paper.

The complete picture of the autumn application cannot be given until next season, when we can estimate the recovery of the different treatments.

CONCLUSIONS

It would appear from this work that MCPB can be used with safety at 2lb. per acre on a mixed sward, but there is doubt whether more than 1lb. is useful on peas. There is probably little advantage in using the butyl ester of MCPB. 2,4-DB would appear to have some superiority over MCPB in the control of weeds, and workers in Britain find it even less phytotoxic to white clover than MCPB. (3, 4, 5, 6). 2,4-DB is useful for weed control in seedling lucerne, but should not be applied during frosty weather to certain varieties.

Finally, I would like to acknowledge the assistance and encouragement I have received from my colleagues and the staff at Invermay, without whose teamwork such a series of trials could not be carried out. I am also grateful to Messrs Irvine and Stevenson and the University of Otago Home Science School for their assistance in processing and tasting of pea samples.

TABLE 1—ANALYSIS OF PEA YIELDS

	lb./acre unshelled	Difference	Shelled pea wt. as % of total	lb./acre shelled
Dinoseb (amine)	15,240	5,880†	47·61%	7,260
Sodium salt MCPB 1lb.	12,460	2,800†	47·07%	5,870
Sodium salt MCPB 2lb.	12,380	2,720†	47·08%	5,830
Control	9,660	—	49·57%	4,790
Sig. diffs. 5%* = 1,470				
1%† = 2,040				
C. of V. = 9·6%				

TABLE 2—PALATABILITY OF SPRAYED PEAS

Flavour	Dinoseb	Unblanched			Dinoseb	Blanched		
		1lb.	2lb.	Control		1lb.	2lb.	Control
1. Desirable	8	8	5	7	10	7	2	7
2. Slightly undesirable, some off flavour	5	6	8	6	4	6	10	6
3. Undesirable, strong off flavour	1	—	1	1	—	1	2	1

TABLE 3—PERCENTAGE CONTROL OF DOCKS

	3 weeks after spraying	3 months after spraying	3 weeks after second application
Potassium salt MCPA	75	20	60
Ethyl ester 2,4-D	75	35	70
Sodium salt MCPB	40	15	25
Butyl ester MCPB	35	25	65
Sodium salt 2,4-DB	60	50	70
Butyl ester 2,4-DB	70	60	80

TABLE 4—LUCERNE YIELDS (LB./ACRE DRY MATTER)

	1st cut	2nd cut	Total
Dinoseb (amine)	2170	1609	3778
Sodium salt 2,4-DB 1lb.	2146	1666	3812
Sodium salt 2,4-DB 2lb.	2146	1633	3779
Control	2290	1676	3966
Sig. diffs. 5%	N.S.	N.S.	N.S.
C. of V.	6·0%	4·4%	4·3%

TABLE 5—EFFECT OF 2,4-DB AND FROST ON LUCERNE VARIETIES

Variety	Frost damage at 8/4/57	Hormone damage at 3/5/57	No. of replicates	
N.Z. Certified	2·25	2·5		
Du Puits	2·0	3·0		
Socheville	2·5	1·75		
Hunter River	3·5	8·25		
Provence	2·75	3·75		
Rhizoma	1·25	1·0		
M. Glutinosa	1·5	1·5		
Gen. San Martin	3	8		
Saladina	3	8		
Argentina	3	8		
			Mean of four replicates	Correlation coefficient r = ·86 P = 0·05
			Only one replicate sown	Not analysed

0 = No damage. 10 = most damage.

TABLE 6—EFFECT OF GAMMA BUTYRICS ON THE YIELD OF SPECIES

Cut No. 1	Total yield	Grass	Red clover	White clover	Weeds
Sodium salt MCPB 2lb.	810	680	50	50	30
Butyl ester MCPB 2lb.	760	620	55	30	55
Control	860	690	85	25	60
Cut No. 2					
Sodium salt MCPB 2lb.	1180	840	200	120	20
Butyl ester MCPB 2lb.	1170	645	360	140	25
Control	1250	730	360	150	10

REFERENCES

1. Anderson, S (1954). Proc. Brit. Weed Control Conference.
2. Rademacher, B. (1954). Proc. Brit. Weed Control Conference.
3. Elliot, J. G. (1956). Brit. Weed Control Conference.
4. Deakins and Ormrod (1956). Brit. Weed Control Conference.
5. Elliott, Green and Evans (1956). Brit. Weed Control Conference.
6. Fletcher and Raymond. Nature, Vol. 178, No. 4525, 21 July 1956.

DISCUSSION

Q.—Have you had any trials comparing low rates of phenoxyacetics with high rates of phenoxybutyrics?

A.—I consider low rates of MCPA would give as good control as high rates of MCPB. I think 2,4-DB will be a more useful material than MCPB.

Comments: At Rukuhia we compared $\frac{1}{2}$ lb., 1lb., and 2lb. of potassium salt of MCPA with 1lb., 2lb., and 4lb. of sodium salt of MCPB. Mowing cuts showed the depressing effect on white clover of 4lb. of MCPB to be less than $\frac{1}{2}$ lb. of MCPA. This effect lasted from spring until late summer.

Comment: I do not think you get any corresponding increase in kill with high rates of butyrics. One and a half pounds may kill as many weeds as 8lb.

Q.—Has any work been done on breeding to increase the resistance of pasture plants to weedkillers?

A.—Plant breeders have a good knowledge of the inheritance of enzyme transmission in clovers. This is a project Grasslands Division, D.S.I.R., Palmerston North, has in mind.

Q.—Were not the rates of 1lb. acid equivalent per acre on docks rather low for such a tough weed?

A.—Possibly a little low, but we wanted to keep rates uniform. The important point established was that 1lb. of 2,4-DB was better than 1lb. of 2,4-D.