On this evaluation it is obvious that if the democratic test of the greatest good for the greatest number is applied, there is a poor case for the relatively few farmers who will benefit from the use of blight jeopardising the soil and water conservation service manuka performs for the whole community. The attempted control of manuka by blight is not consistent with the over-all demands of soil and water conservation, that is, a balanced use of protective and productive vegetation to assure maximum permanent production from as much land as possible.

As so much of our unfarmable land requires protection, it must be fully realised that manuka is the miracle plant of marginal land.

Forester's View

F. W. FOSTER, Ministry of Works, Wellington.

THAT a forester should be taking part in this symposium is somewhat anomalous, for manuka is neither a forest weed nor a land-use weed. Therefore it is not a national weed.

Many agriculturists (though not all) consider it a noxious weed, but it has never been so proclaimed. In an official report of 1909 entitled "Forestry in New Zealand" a Commissioner of Lands expressed the following opinion on manuka: "Should be in the category of noxious weeds." He was not a forester, and his opinion was not acted upon, which is understandable in view of the large areas of infertile beech forest land opened up in the district.

It is necessary to note the two common kinds of manuka: *Leptospermum scoparium*, manuka, red manuka, or scrub manuka; and *L. ericoides*, white manuka or tree manuka. The latter grows to twice the size of the former in both height and diameter.

Cheeseman's "Manual" gives their distribution as abundant throughout the North and South Islands and ascending to 3000ft. and 4500ft. altitude respectively. Kirk in his "Forest Flora" rightly says that white manuka is rare in South Otago and Southland. He says of red manuka: "The most common plant in the colony, and found in great abundance in every district." That was in 1889, when refrigeration and consequent increased prices for meat were beginning to extend settlement more and more on to hill country. Thus there was an abundance of manuka shrubland before the reversion of settled hills began on any appreciable scale.

Probably manuka was not so widespread or abundant in 1840 as at present, though the Maori used to practise a form of shifting cultivation necessitating clearing forest patches which were later abandoned. Manuka would colonise these or their margins. The Maori, too, used to burn. Hochstetter, writing in 1867 of the kauri region, said: "What formerly had been employed in the war of cannibal tribes, in a stratagem to burn out the enemy, is done now for the sake of money." Volcanic eruptions long ago would be colonised later by manuka, as happened in recent times around Mt. Tarawera. After settlement began, surveyors', gumdiggers' and miners' fires, as well as those of runholders and other settlers, all favoured the extension of manuka.

Doubt has been expressed as to whether manuka actually does function as a nurse for forest regeneration. It most certainly does.

Cheeseman writes that the heath-like red manuka vegetation often stretching for many miles "will, if left alone, gradually develop into a taller growth until finally, by the appearance of young forest trees among the manuka, the beginning of a mixed-forest growth is seen, which ultimately supersedes the manuka." The same writer gave further detail in a paper read to the N.Z. Forestry League in 1917. "What happens," he says, "is this: The manuka, as it gradually increases in height, becomes a little more open at the base and forms a sheltered nursery for the seedlings of many kinds of trees, derived, no doubt, from seeds conveyed by the wind or otherwise from an adjacent forest. As years go by, the manuka approaches its full height, and at the same time becomes more open at the base; the growth of the trees is accelerated, so that they ultimately overtop the manuka, finally becoming a young and vigorous forest. This formation of forest through a natural process can be seen in numerous localities throughout the whole of the North Island." He added that the value of manuka as a nurse for young forest is an indisputable fact.

It should be noted that the mature forest need not be actually adjacent to the scrub, for seed is commonly dropped by birds which perch on or nest in the manuka.

Dr. Cockayne also described the succession of forest on manuka shrubland if left unharmed by burning or cutting. (We can now add "or by manuka 'blight'.") The process must be a natural one, though foresters may one day develop a silvicultural technique to hasten the process. I personally attempted this on one occasion by removing 75 per cent. of the manuka, but I was too early or too severe, for the seedling mortality 3 years later far outweighed the recruitment from additional seedlings which had germinated in the meantime.

The Forest Service has more recently successfully liberated kauri at a later stage in its growth, and moreover has actually broadcast white manuka seed in order to establish a nurse for kauri regeneration. White manuka is the better nurse, but its occurrence is often sporadic, and in any event red manuka, generally abundantly present, serves as a nurse for the white manuka.

The following table shows what happens at the margin of a kauri forest when manuka is not interfered with. The counts and measurements were made by the writer in Waipoua Forest some years ago.

Forest Seedlings and Trees per Acre under Manuka

Age of Vegetation.	Kauri.	Other Conifers.	Broad- leaved.	Total.
Seedlings (to 3ft. tall)				
Not over 8 years ..	1550	100	50	1,700 Manuka & fern 2-3 feet.
About 20 years ..	8300	3100	3350	14,750 Manuka 6-15ft.
Not over 37 years ..	7050	7300	6700	21,050 Manuka 12-25ft.
Trees (4in. diameter and over)				
About 70 years ..	247	67	190	504 Height of Stand 55 feet.
About 150 years ..	189	57	193	439 Height of Stand 85 feet.

On the 20-year area manuka were under 9in. apart or nearly 80,000 stems per acre. In the 70-year area there were 125 white manuka and 10 red manuka; and of these 48 per cent. of the former but only 20 per cent. of the latter were still of vigorous growth. The white manuka was up to 50ft. tall and 15in. diameter, and the red manuka about 20ft. tall and 8in. diameter. Incidentally, the kauri diameters here were 8in. average and 18in. maximum.

In the 150-year stand there were 17 manuka, all white, the red having long since been suppressed and died. They were up to 80ft. tall and 16in. diameter, but were fast going out, the crowns being a mere tuft about 3ft. across. (Kauri here were 15th. average diameter to 37in. maximum.) On this acre there were perhaps 20 dead manuka, the trunks rotted off at up to about 40ft. above the ground. The remarkable thing is that there should be any manuka still alive and towering to such a height.

The inescapable conclusion is that the manuka has protected and drawn up the kauri for well over a century, though its job is now completed.

The writer has observed forest regenerating in manuka in numerous localities extending from Hokianga to Otautau. Abundant regeneration of rimu was seen near Taheke in the north, in Taranaki, and elsewhere. Indeed rimu comes in very much better under manuka than under mature forest. Near Katikati I have seen an odd mixture of kauri, hard beech, and insignis pine, all regenerating together in manuka. Another forest species that develops well in manuka, and eventually supersedes it, is rewarewa or honeysuckle, as on a large area near Kotemaori, and at Wainui-o-mata, Mt. Tarawera, and elsewhere. Beech regenerates in manuka in the South Island and in exotic pines at Hokianga, Silverdale, Taupo, Nelson and North Canterbury.

Lest it be gathered that the forester is concerned with the bad effect of manuka blight on regeneration of timber trees alone, let me hasten to say that of even greater significance is the effect on young or potential protection as distinct from production forests. Manuka is an excellent pioneer in colonising upland infertile areas damaged by fire, slipping, or wind erosion. Forest species come in, the land is useless for any purpose but conservation, and correct land use demands that it be vegetated in protection forest. Such a forest, however, will not develop where the manuka is removed at the wrong stage. It will develop satisfactorily only when Nature is allowed to complete the succession. Where the manuka is not sufficiently advanced, destroying it will possibly so interfere with the development of succession to high protection forest as to prevent development into a forest at all, or at least retard it for a long period. If this happens and the ground is not well covered in an effective vegetation, erosion of various kinds is likely to occur and flooding to be accentuated.

In the main we should be concerned about extensive protection forest areas above the hill farms. But on hill farms themselves there are often sites too steep, unstable, or infertile to carry pasture safely and economically, and such sites should be clothed in the vegetation which will cover them most effectively. Manuka is one of the plant associations which in many instances is actually doing this job, and it is in the farmer's interest to conserve it. The areas in question could be planted in exotic trees, but are often too steep and inaccessible for timber-getting at maturity, besides being expensive to plant.

While this desirability or necessity of growing forest, for the public well-being as well as the farmer's, on land unsuitable for pasture is being dealt with, certain agricultural and pastoral statistics might be briefly referred to. Agriculturists, I imagine, study the statistics of improved land, and I hope to be forgiven for looking at the unimproved land areas. The occupied area in scrub, fern and second growth increased by nearly 2 per cent. between 1922 and 1932, 9½ per cent. in 1932-1942, and 23½ per cent. in 1942-1952. Between 1942 and 1952, when the total unimproved and total occupied areas remained fairly constant, the percentage of total occupied land which comprised scrub, etc., rose from 10.1 per cent. to 12.4 per cent. The tussock-danthonia percentages dropped from 32.3 per cent. to 30.7 per cent.