Research Note

RAGWORT CONTROL WITH PICLORAM

P. G. COLES

Ivon Watkins-Dow Ltd., New Plymouth

Summary

Results from field trials for the control of ragwort (*Senecio jacobaea*) are presented. Treatments containing picloram either alone as a granular formulation or in combination with 2,4-D as spray treatments have consistently proved more reliable than the butoxy-ethanol ester of 2,4-D. Broadcast applications of 1 oz picloram plus 4 oz 2,4-D per acre; spot applications of a concentration containing 2.25 oz picloram plus 9 oz 2,4-D per 100 gal and broadcast granular treatments of 8 oz picloram per acre all gave effective control of ragwort in the advanced rosette stage. Higher rates appear necessary to control ragwort in the flowering stage but further studies are necessary to establish minimum effective rates of both granular and spray formulations at the various stages of ragwort growth. The effect of picloram/2,4-D mixtures on ragwort seed viability is discussed.

INTRODUCTION

RAGWORT (*Senecio jacobaea*) is a native of Europe and Asia, and was introduced to New Zealand last century, being first recorded in the South Island in 1874. It has now become a major weed in both islands, even on quite fertile soils, but especially in dairy pastures on the light pumice soils of the central North Island, where the pasture cover can be more readily opened up in dry summers. It was first declared a noxious weed in 1900 and, because of its alkaloid content, it has also been recognized as a poisonous plant, especially to cattle.

Ragwort is usually considered a biennial but under certain growth conditions it can become a perennial with the ability to make prolific root and crown regrowth (Poole and Cairns, 1940). Until the introduction of 2,4-D, control measures were confined mainly to grubbing, hand pulling, and spot treatment with sodium chlorate. Owing to the strong regrowth characteristic of ragwort, however, variable results have been obtained with these methods. More recently, 2,4-D either as the ester, amine or the sodium salt formulation has been the preferred treatment, with the esters showing superior activity to the amine and salt formulations. Up to the advanced rosette stage of growth, control with 2,4-D has been reasonably satisfactory. With 2,4-D treatments of older plants, which have developed mature crowns with a reduced leaf ratio, results have been variable, with regrowth frequently appearing.

In 1964, picloram/2,4-D as an amine salt formulation became commercially available. This product was recommended for spot treatment of annual and perennial broadleaf pasture weeds among which ragwort proved highly susceptible. Picloram has proved more readily translocated than 2,4-D and, because of its residual properties, has shown outstanding control of regrowth, proving complementary to the quicker acting 2,4-D. In 1965 a granule formulation of picloram became available and this, too, showed excellent activity against ragwort. This paper presents the results of field trials to control ragwort with these picloram formulations.

EXPERIMENTAL

For replicated small plot boom applications, an Oxford Precision Sprayer was used to apply 32 gal per acre to $\frac{1}{200}$ th acre plots (31 ft $\times$ 7 ft) at 30 lb/sq.in. and 3 mph. Granule applications were broadcast by hand to 16 sq.yd plots using an inert filler to ensure more even coverage of the plots.

Spot spray applications were applied with a hand wand using a standard engine-driven pump unit at 80 lb/sq.in. to give a light but thorough foliage wetting. All rates of application are expressed as active ingredient either on a per acre or on a per 100 gal spray mixture basis. Assessments were carried out on a 0 to 10 numerical scale by two or more independent observers. (0 = no effect; 10 = complete control). The 2,4-D used as the standard treatment was as the butoxy ethanol ester, while the picloram/2,4-D mixture was as an amine salt formulation. The picloram granules were based on an absorbent attapulgite clay core.

RESULTS AND DISCUSSION

BROADCAST APPLICATIONS

Trial 1: November 1965, Taranaki (Free-draining brown loam)

The ragwort plants were in the advanced "cabbage" to flower bud stage, ranging in height from 12 to 42 in. A light drizzle fell following applications, but the rain was insufficient to cause run-off from the leaf. At this rather advanced stage of treatment, picloram/2,4-D at 1 + 4 oz was about equivalent in control to the standard 2,4-D ester at 20.8 oz, both giving marginal final kill (Table 1).

An acceptable final level of control was obtained with the higher treatment rates, but complete kill of all plants was still not achieved from this late stage broadcast spray treatment.

Trial 2: May 1966, Bay of Plenty (Light pumice ash)

At this late autumn application, the ragwort was in an early "cabbage" stage, prior to appearance of the flower stalk, the plants being from 3 to 8 in. in height. The evaluation made after 4 months indicated that picloram/2,4-D at 2 + 8 oz was necessary for complete kill, while at 1 + 4 oz an acceptable if not quite complete control was achieved (Table 1).

TABLE 1: INITIAL AND FINAL CONTROL OF RAGWORT IN TWO TRIALS FROM PICLORAM/2,4-D BROADCAST TREATMENTS

Treatment (oz/acre)		7	Control			
			Trial 1	Elapsed Time (weeks)		
			11	8	32	
2,4-D ester (standard)	20.8		2.4	6.8	0.5	8.0
Picloram 0.5/2,4-D 2			—	—	1.7	7.0
Picloram 1/2,4-D 4			2.9	6.3	3.3	9.0
Picloram 2/2,4-D 8			4.4	8.2	5.7	10
Picloram 3/2,4-D 12			6.1	8.9	—	—
Picloram 4/2,4-D 16			—	—	7.3	10
Untreated control			0	0	0	0

This is in general agreement with earlier results by Schmidl (1965). Both the 0.5 + 2 oz rate of picloram/2,4-D and the standard treatment of 2,4-D ester at 20.8 oz gave less than acceptable results. It was noted that clover protected under the rosette of leaves survived the spray treatment and grass regeneration into the ragwort areas was rapid and complete following the death of the ragwort.

Trial 3: December 1966, Taranaki (Egmont volcanic ash)

On this site 0.5 in. of rain fell within 24 hours, starting less than one hour after treatment. Ragwort was in a vegetative to early bud stage, varying in height from 4 to 30 in.

No foliage brown-off or control was noted three months following treatment except that the 4 + 16 oz picloram/2,4-D rate had suppressed flower-head formation. It was concluded that the heavy rain so soon after application had reduced the uptake of the chemicals to a sublethal level, as the spray droplets would have not even dried on the foliage.

Trial 4: October, 1965, Taranaki (Free-draining brown loam)

A spot spray treatment applying about 50 gal per wetted acre was made to actively growing ragwort in the seedling to advanced "cabbage" stage of growth. At this stage of plant growth, the standard treatment of 2,4-D ester and all picloram/2,4-D rates above 2 oz picloram per 100 gal gave complete final kill of ragwort (Table 2). At 1.12 + 4.5 oz picloram/2,4-D, slightly inferior, but acceptable final control was achieved. It was noted, however, that 12 weeks after treatment seedling regrowth was present in the 2,4-D ester treated plots, while only trace seedling regrowth was noted in the plots sprayed with 1.12 + 4.5 oz picloram/2,4-D concentration. The thorough wetting obtained using spot treatment applications of picloram/2,4-D showed more rapid plant reaction than with the broadcast low volume treatments.

TABLE 2: SPOT TREATMENT CONTROL OF RAGWORT WITH PICLORAM/2,4-D

Treatment (oz/100 gal)			Control Elapsed Time (weeks)	
			5	12
2,4-D (ester standard) 48			7.5	10
Picloram 1.12/2,4-D 4.5			8.5	9.0
Picloram 2.25/2,4-D 9.0			8.5	10
Picloram 4.5/2,4-D 17.8			9.2	10
Picloram 6.7/2,4-D 26.9			9.2	10
Untreated control			0	0

Trial 5: December 1966, Taranaki (Egmont volcanic ash)

The ragwort plants were in a vegetative to early bud stage and varied in height from 4 to 30 in.

Complete root kill was not obtained from any treatment in this broadcast granule application (Table 3). 12 oz picloram proved to be a marginal dose for older plants with massive crown and mature root systems. It was noted after 7 weeks that only seedling plants had been

controlled but after more than 12 weeks some of the older flowering stage plants had also succumbed.

With ragwort at a slightly earlier stage, picloram granules down to 8 oz broadcast application have given eradication (Moffat, 1966).

TABLE 3: RAGWORT CONTROL WITH PICLORAM GRANULES

<i>Treatment (oz/acre)</i>	<i>Elapsed Time—3 months</i>	
	<i>Top Kill</i>	<i>Root Kill</i>
Picloram 12	5.8	5.5
Picloram 24	9.6	8.9
Picloram 48	9.6	9.5
Untreated control	0	0

Farmer applications of picloram granules are normally made as spot treatments at the rate of a half teaspoon per plant (approximately 0.04 gram picloram) placed in the centre of the well-developed rosette. Such field-scale treatments have consistently given complete ragwort control regardless of the season of application.

Trial 6: January 1967, Taranaki

The purpose of this trial was to assess the viability of seed harvested from ragwort sprayed in full flower.

Results obtained indicate that picloram/2,4-D at 4 + 16 oz can markedly reduce the percentage germination of seed (Table 4), and also result in production of a large percentage of abnormal seed which normally dies a few days after germination. At this late stage treatment lower rates of picloram/2,4-D showed little effect on seed viability, though the same treatment rates applied at pre-flower stage would prevent the formation of flower heads.

TABLE 4: RESULTS OF SEED VIABILITY TESTS FROM PICLORAM TREATED RAGWORT IN FULL FLOWER

<i>Treatment (oz/acre)</i>	<i>% Germination</i>	<i>% Abnormal Seed</i>
2,4-D ester 32.0	20	17
Picloram 1/2,4-D 4	47	12
Picloram 2/2,4-D 8	31	12
Picloram 3/2,4-D 12	26	10
Picloram 4/2,4-D 16	9	27
Untreated control	33	16

CONCLUSIONS

Picloram/2,4-D treatments of ragwort up to the advanced rosette stage of growth have been consistently more reliable than the previous standard 2,4-D ester treatment. In comparison with 2,4-D ester, the picloram/2,4-D mixture has also given much superior kills of mature ragwort in the advanced rosette to full flowering stage of growth. Crown regrowth is normally prevented.

Future studies should attempt to establish the minimum dosages of both picloram/2,4-D spray and picloram granule formulations required for complete ragwort control at varying stages of growth particularly as spot applications.

More critical seed viability studies should be carried out to determine the effective treatment stage and dosage rate for the elimination of seed production in flowering ragwort.

ACKNOWLEDGEMENTS

Grateful thanks are due to Dr G. W. Mason for assistance in preparation of this paper, to various members of the Research and Development Department, Ivon Watkins-Dow Ltd. for assistance with the trial work; to farmers who maintained the trial sites for continued evaluation, and to the Department of Agriculture, Seed Testing Station, Palmerston North, for seed viability results.

REFERENCES

- Moffat, R. W., 1966: *Proc. 19th N.Z. Weed & Pest Control Conf.*: 93.
Poole, A. L.; Cairns, D., 1940: *Dept Sci. ind. Res. Bull.* 82.
Schmidl, L., 1965: *Aust. Weeds Conf.*, 1: 5.