SELECTIVE WEED CONTROL IN POTATOES

L. J. MATTHEWS

Research Division, Department of Agriculture, Hamilton

Summary

An inadequate number of trials have indicated the value of herbicides for potato weed control applied just prior to shoots breaking the surface of full ridges made at the time of planting. Results based upon potato trials alone are not finite enough to list herbicides in order of efficiency, but selective foliage-active, residual-type herbicides appear to be the most suitable materials. No cultivation comparisons were made, all yields being compared with hand-weeded control. In all trials, differences of less than 1 ton per acre were not significant, hence yields are not quoted.

INTRODUCTION

REALIZATION that post-emergence cultivations spread virus diseases, cause loss of soil moisture, sometimes loss of yield, eventual deterioration of soil structure, and foster further seed germination has increased the desire to employ selective herbicides. Traditionally, potatoes have been regarded as a cleaning crop and the art of cultivation for weed control alone has been developed to a high degree.

Reliance may be placed on herbicides to supplement cultivation by keeping the crop clean until emergence, or after crop emergence, or throughout the duration of the crop. In this latter case, the crop is fully ridged at the time of planting and no further cultivation is practised. Ten statistical yield trials have been conducted over the past three years comparing a range of herbicides with hand-weeded control (ridged at the time of planting and hand-weeded). Reference is made to a number of previous trials.

RESULTS

All rates are expressed in terms of active ingredient per acre.

STAGE OF GROWTH

In three trials the same herbicide was applied at more than one stage of growth. Propanil 2 lb in one trial yielded significantly better at 1% applied at pre-emergence than at post-emergence. There was no difference in the 1 lb rate, both being significantly worse than hand-weeded control. In two other trials, rates of 1, 2 and 4 lb applied at 50% emergence, 2 to 6 in. stage, 6 to 12 in. high, and at pre-flowering, did not differ significantly from hand-weeded control. For simazine and prometryne, yield differences were not significant for 5% emergence and the 2 to 6 in. stage, though depression was greater at the advanced stage of growth. In all other cases, treatments were applied at the first sign of emergence.

SOIL TYPE

On a clay loam, simazine did not reduce total yield but increased proportion of seed and pig potatoes (below 2 oz in weight). On silt loam in two trials, yields were depressed. On a sandy loam, yields were depressed by 15 and 13% but differences were not

significant. Only 1.1 in. of rain fell in the first month of application. Prometryne gave similar results only at the 2 lb rate applied at time of planting on a clay loam. All other treatments did not differ significantly from hand-weeded control.

WEATHER CONDITIONS

Heavy rain, particularly after a dry period, is more likely to cause damage, as many potato roots are nearer the surface. As indicated under "Soil Type", heavy rain, especially on soils low in organic matter, make potatoes more susceptible to residual-type herbicides. Given adequate rain, a relatively insoluble material such as simazine can reduce yields even on a clay soil high in organic matter. Monolinuron, because of its greater solubility, shows better residual weed control under dry conditions than linuron, simazine, prometryne and ametryne. The speed of action of most herbicides is more rapid under warm, moist conditions.

HERBICIDES

In the trials the main types of herbicides considered for potato weed control were as follows:

(a) Activity through Foliage

In previous trials, MCPB, MCPA, 2,4-DB and 2,4-D caused significant depression over hand-weeded control and their use was justified only if susceptible perennial weeds dominated. Propanil and the allied material BV207 have not given a wide enough range of weed control over all conditions to be useful unless supplemented by cultivation. Applied before potato foliage is present, diquat and paraquat have given a wide range of weed control and would be useful where cultivation after emergence is the usual practice, such as for heavy soils that encrust. The same applies to phenols alone or in conjunction with aromatics, but here there is a short residual control.

(b) Activity through Soil

Previous work with diuron showed results were too variable to recommend this herbicide. Results with simazine at 1 lb and 2 lb have indicated the same. Even if allowance is made for soil type, other variables, such as rainfall, are unpredictable and excessive damage may result.

(c) Activity through Foliage and Soil

To date, these materials have given the most consistent results. The methoxy-ureas, linuron, monolinuron and metobromuron and several of the 2-methyl-mercapto substituted triazines have all given favourable results. These compounds have sufficient foliage activity on a broad spectrum of weeds to warrant use. In a season where successive strikes of weeds occur, their residual action ensures prolonged control provided resistant species are not dominant. In several trials their effectiveness has been such that a pre-harvest desiccant would not be required.

RATES OF APPLICATION

Under warm conditions and when the weeds are in the seedling stage of growth, rates of 4 to 8 oz of the methoxy-ureas and 2-

methyl-mercapto substituted triazines are adequate for susceptible species. These rates possibly do not have sufficient persistence in light sandy soils or adequate herbicidal effect in heavy clay soils. Rates of 1 lb appear optimum for lighter soils and should persist under most conditions until the crop meets over the row. For heavier soils, rates higher than the 1 lb may ensure more adequate weed control but could result in crop damage in a wet season. Under very wet conditions, even a 2 lb rate of prometryne failed to give adequate weed control, and crop damage was significant.

WEED SPECIES CONTROL

The presence or absence of weed species dictates to some degree the usefulness of foliage-residual herbicides.

Annual Weeds

Fumitory (*Fumaria officinalis*), *Galium* species, *Veronica* species and *Stachys arvensis* show both pre-emergence and post-emergence resistance to the methoxy-urea compounds. *Anthemis cotula*, *Matricaria* species and *Polygonum aviculare* show post-emergence resistance, particularly to monolinuron and metobromuron. The addition of a wetting agent to linuron has improved the foliage activity to such an extent that satisfactory post-emergence control of the more resistant weeds may be obtained. Monolinuron is superior to linuron in control of annual grasses.

Cruciferous weeds, *Ranunculus* species, *Galium* species (prometryne in particular), *Matricaria* species, *Polygonum aviculare* and some grasses show high resistance to the 2-methoxy-mercapto substituted triazines and *Solanum* species are suspect.

Perennial Weeds

The presence of *Cirsium arvense* necessitates the employment of cultivation or MCPB and MCPA. The same applies to rhizomatous grasses — it being necessary to employ trichloroacetic acid.

Stage of Growth of Weeds

This, to some extent, is dictated by the time the crop takes to emerge. Ideally, the crop should be within a day or so of breaking the surface when the weeds are in the seedling stage of growth. Beyond the seedling stage of growth, the more difficult to kill weeds show higher tolerance. This is true especially for metobromuron and monolinuron which do not have the same foliage activity as linuron at equivalent rates. This also applies to prometryne on species controlled by both groups of chemicals — prometryne having less activity than linuron.

Varieties

In three trials a number of varieties were included. In no instance was the varietal response significant. The varieties included Sebago, Glen Ilam, Auckland Short Top, Ilam Hardy, and Rua.

TABLE, SEED AND PIG POTATOES

Potatoes are difficult to kill but are easily damaged. Even where lack of visual damage is evident on the foliage, yields or grading may be reduced. Reduction in grading quality — i.e., higher proportion of seed and pig potatoes — usually signifies less damage has occurred than when total yields are reduced.

Potatoes are a very variable crop and in a number of trials reported the experimental error has been so high that yield differ-

ences have not been significant, although large differences in results are apparent — *e.g.*, simazine 2 lb depressed yields by 15% but as this depression was less than 5.3 tons to the acre the figure for minimum difference for significance results were not significant from hand-weeded control or propanil 2 lb which gave the highest yield. The trial was replicated four times and few, if any, weeds, were present.

RESIDUAL DAMAGE TO OTHER CROPS

This aspect has not been followed specifically in trials. There have been no adverse reports on the methoxy-urea compound or the 2-methyl-mercapto substituted triazines but suspected damage to cucurbits has followed the use of simazine/prometryne mixtures.

DISCUSSION

Herbicides for potato weed control have been evaluated in comparison with hand-weeded control, not unweeded control receiving no cultivation. It is concluded that hand-weeding would be satisfactory as conventional cultivation except on heavy soils that encrust. In some cases, hand-weeding may produce higher yields than cultivation through lack of crop damage, hence herbicides that produce yields equivalent to hand-weeded control require extensive evaluation.

Potato yields are variable and often large depressions do not reach significance. Unless differences of less than 1 ton per acre are significant, it is difficult to evaluate the role of herbicides for potato weed control. However, the trials show that at least two groups of herbicides are promising, namely, the methoxy-ureas and the 2-methyl-mercapto substituted triazines. Each group has advantages and disadvantages, though the disadvantages are small in comparison with previous type herbicides. New Zealand experience would tend to confirm English work (Pheiffer *et al.*, 1964) that simazine is not a reliable chemical to add to the 2-methyl-mercapto substituted triazines. Under dry conditions, the effects are largely nullified, and under wet conditions damage may be apparent in total yields or in grading. The range of weeds controlled is very similar.

Prometryne has given excellent control of *Spergula arvensis* and *Fumaria* species even at the rate of 6 oz. Control of the former species is not always positive with methoxy-ureas and control of the latter is obtained only with 1 to 2 lb rate plus wetting agent. Similarly, the methoxy-ureas are more active on cruciferous weeds and *Solanum nigrum*. A mixture of the two groups would, no doubt, give a broader spectrum of weed control.

Much further information is required on effects of soil type and rainfall. Potatoes, at present, are grown on soil types from light sands to heavy clays and rainfall varies from 25 to 60 in. depending on the district. It is not feasible to determine all variables by trials, hence the final use of foliage-residual herbicides will depend on farmer understanding.

ACKNOWLEDGEMENTS

Thanks to due to P. Bramley, A. Thompson, and F. A. Meeklah for conducting field experiments and to the Superintendent, Biometrics Section, for statistical analysis.

REFERENCE

Pheiffer, R. K., *et al.*, 1964: *Proc. 7th Brit. Weed Control Conf.*: 411.