WEED CONTROL IN NEWLY TRANSPLANTED ASPARAGUS

A. RAHMAN, A. THOMPSON and P. SANDERS

Ruakura Soil and Plant Research Station, MAF, Hamilton

SUMMARY

Five trials were conducted over a 2 year period to investigate general herbicides for weed control in newly transplanted asparagus (*Asparagus officinalis*). Simazine 1.5-2.0 kg/ha, diuron 1.5-3.0 kg/ha and bromacil 0.75-2.0 kg/ha applied pre-emergence showed useful broad spectrum activity against weeds with apparently good crop tolerance. Good residual weed control was achieved from bromacil for 4-6 months while simazine and diuron treatments lasted for only 2-4 months. Post-emergence control of weeds was best obtained by a mixture of metribuzin 0.4-0.5 kg/ha plus diuron 2.0-2.5 kg/ha as a directed spray to minimise crop damage.

INTRODUCTION

Because the long term marketing prospects for asparagus look bright, it has created intense interest among Waikato farmers contemplating diversification into horticulture. This is tempered a little by the knowledge that in Hawkes Bay, where asparagus has traditionally been grown, yields have declined and as a result it is being replaced by more profitable crops (Dawbin 1980). Thus, the area planted in this crop has risen rapidly to an estimated 350 ha in Waikato and 300 ha in the Bay of Plenty region by the end of 1980 (S.J. Franklin pers comm) and several nurseries have been established for plantings in the coming season.

Weed control is a major problem of asparagus as weeds not only compete with the crop but are also known to restrict the air movement through the crop, resulting in higher humidity which could increase the incidence of the fungus *Stemphylium*. Weed seed set in the crop can also aggravate the weed problem in the future. In addition to the weeds encountered in the traditional areas of asparagus growing, growers in Waikato and Bay of Plenty have to contend with rampant growth of warm zone grasses such as paspalum (*Paspalum dilatatum*), summer grass (*Digitaria sanguinalis*), smooth witchgrass (*Panicum dichotomiflorum*) and barnyard grass (*Echinochloa crus-galli*) (Franklin 1980). Limited experience by local growers indicates that weed control recommendations developed for other areas will require considerable modifications to meet the differences in soil, climate, and the weed spectrum.

The work reported in this paper summarises preliminary results of five trials conducted over the past two growing seasons in newly transplanted asparagus.

METHOD AND MATERIALS

Four trials were conducted to evaluate herbicides applied pre-emergence (pre-E) and one trial post-emergence (post-E) between 1979 and 1981. The soil type was Horotiu sandy loam except Trial 2 which was Te Kowhai silt loam. Further details of the trial sites are given in Table 1. The plots were 8-13 m long and 2-3 m wide in which asparagus crowns cv. Mary Washington 500W had been transplanted at 1 m between rows (cv. Limbras at 2 m row width in Trial 4) and 25-45 cm apart within rows. Treatments were arranged in a randomised block design with four replicates. All herbicides were applied in 350 litres /ha at 200 kPa.

Eleven herbicides alone or in mixtures were investigated for pre-E weed control and were compared with untreated control plots. When treatments broke down, inter-row

Crops

TABLE 1: Details of trials.

Trial No.	Location	Organic C (%)	pH	Rainfall* (mm)	Planting date	Spraying date
1	Hamilton	3.6	5.9	115	10.9.79	14.9.79
2	Te Kowhai	5.9	5.6	103	24.9.79	27.9.79
3	Rukuhia	8.8	5.4	112	15.8.80	22.8.80
4	Tamahere	5.4	5.8	44	30.9.80	10.11.80

Trials 1-3 pre-emergence, 4 post-emergence

* during 4 weeks after spraying.

cultivation was carried out in Trials 1 and 2 and metribuzin 0.4 kg/ha was applied as a directed spray in Trial 3.

In Trial 4, several herbicides alone or in combination were investigated for post-E use. Except for two treatments which were sprayed overall, the herbicides were applied as a directed spray using a hand-held boom between the 2 m rows. In this way only the lower portion of the stem and any ferns close to the ground received the chemical. Weed growth at the time of treatment was up to 10 cm in height, principally of redroot (*Amaranthus powellii*) and black nightshade (*Solanum nigrum*).

Assessments of the weeds were made by visual scoring, seedling counts and dry matter production, while stem counts and visual assessments of fern damage were recorded for the crop.

RESULTS AND DISCUSSION

The main weed species encountered in these trials were redroot, black nightshade, twin cress (*Coronopus didymus*), willow weed (*Polygonum persicaria*), summer grass and poa annua (*Poa annua*). Other weeds present in smaller numbers included prostrate amaranth (*Amaranthus deflexus*), narrow-leaved plantain (*Plantago lanceolata*), annual mouse-ear chickweed (*Cerastium glomeratum*), creeping buttercup (*Ranunculus repens*) and white clover (*Trifolium repens*). At Rukuhia, the trial area was subjected to a very high weed pressure as is indicated by the weed numbers in Table 2. For pre-E weed control, simazine was reasonably effective and was slightly superior to diuron in all trials. Linuron also gave effective weed control, but of slightly shorter duration than diuron. Prometryn tested in Trials 1 and 2 did not provide effective weed control, results of which are not tabulated. Eventually all of these materials were overwhelmed by redroot, summer grass and in Trial 3, by paspalum.

Bromacil, even at the low rate of 0.75 kg/ha was much more effective than the other materials, both in the spectrum of weed species controlled and in persistence. Weed invasion, mainly by summer grass and some redroot started to occur 5-6 months after treatment in Trials 1 and 2. Paspalum was able to establish in Trial 3 after 4-5 months and eventually dominated the trial area. The addition of diuron to bromacil for pre-E use was of little benefit in these trials.

There was no apparent evidence of adverse herbicide effect on the asparagus crop at any site. Significant differences between stem counts or stems/plant were not recorded between various treatments. Slight reductions in numbers of stem counts in Trials 1 and 2 (Table 2), may be due to competition from weeds remaining within the rows, as cultivation was carried out only between the rows after the treatments had broken down.

A separate trial to investigate other possible chemicals for pre-E use was laid down in small plots next to Trial 3 on the same date. Treatments included trifluralin plus linuron, alachlor, hexazinone, pendimethalin and a commercial mixture of atrazine plus metolachlor. The best weed control and longest residual life was from hexazinone and atrazine plus metolachlor, but both these treatments caused some fern damage. Trifluralin plus linuron gave better initial weed control than alachlor, but residual activity was quickly lost and these were no better than any of the treatments in Trial 3. More work is required with some of these treatments to test application rates and

Crops

TABLE 2: Effects of pre-emergence treatments on weed numbers and asparagus stem production.

Treatment Trial No.	(kg/ha)	Weed No./m ²			Asparagus stems/10 m row		
		1	2	3	1	2	3
untreated		74.3*	184.8*	2959*	21.9	34.8	
simazine	0.75	8.7 ab	14.9 b		25.2	36.3	
sinazine	1.5	0.9 cd	1.6 c	54 c	26.7	37.3	
simazine	2.0			66 c			128
diuron	1.0	5.4 bc	21.9 a		23.8	34.8	
diuron	1.5	0.9 cd	4.0 c		28.5	32.1	
diuron	3.0			156 b			144
linuron	0.75	10.9 a	12.7 b		23.1	37.3	
linuron	1.25	3.1 cd	4.6 c		25.4	33.1	
bromacil	0.75	0.4 d	0.0 c		27.9	38.5	
bromacil	1.5	0.0 d	0.0 c	42 c	26.3	38.3	
bromacil	2.0			43 c			142
+ diuron	1.0 + 1.0			53 c	NS	NS	NS
CV%		39.8	27.9	20.3	10.2	9.5	-
days post treatment		54	43	46	66	53	152

NS — no significant differences between treatments; blank spaces indicate treatment not included or data not collected.

* data excluded from statistical analysis.

suitable combinations.

Among the post-E treatments in Trial 4 (Table 3) simazine showed the least crop damage but gave poor weed control, probably because of the presence of some weeds at the time of application. Hexazinone resulted in good control of weeds but crop damage was unacceptable. The best treatment for both weed control and crop tolerance was metribuzin plus diuron as a directed spray. The other two metribuzin mixtures gave results inferior to that of metribuzin plus diuron, especially in residual activity. There was little difference in weed control between the two methods of application, but overall boom spraying caused more crop damage than the directed spray.

TABLE 3: Effects of post-emergence treatments on weeds and transplanted asparagus (Trial 4).

Treatment	Rate (kg/ha)	Weed score †		Crop damage (%)
		21.11.80	15.1.81	
untreated		0	3.0**	0
hexazinone	1.0	8.6	6.7	45
simazine	2.0	1.2	6.7**	1
linuron + metribuzin	0.5 + 0.4	8.0	3.7	20
diuron + metribuzin*	2.0 + 0.4	8.4	4.7	23
diuron + metribuzin	2.0 + 0.4	8.8	5.5	10
diuron + metribuzin	2.5 + 0.4	8.4	7.7	13
methabenzthiazuron + metribuzin*	1.5 + 0.4	7.9	1.2	35
methabenzthiazuron + metribuzin	1.5 + 0.4	8.0	0.5	16
methabenzthiazuron + metribuzin	2.0 + 0.5	8.2	3.5	13
CV%		4.0	20.9	16.4

† 0 = no weed control; 10 = complete weed control

* boom spray overall

** handweeded 12.12.80

Crops

The work reported here has shown some potentially useful herbicides and many of the problems associated with weed control in this crop. Because the economic life expectancy of the crop is at least 10 years, long term effects of herbicides on crop production are very important and need to be examined. On the basis of these short-term trials, bromacil pre-E with its long residual life and broad weed spectrum has the advantage of fewer weed control operations during the season, thus resulting in less soil compaction by heavy equipment and physical damage to the ferns.

Among post-E treatments methribuzin alone or in combination with diuron or methabenzthiazuron appears promising for control of broadleaf weeds, but more work is required on crop tolerance. A number of selective herbicides for control of grasses and cultural methods of weed control in conjunction with chemicals are also being studied.

In the Rukuhia trials where soil organic matter is higher than the other sites and weed pressure was known to be quite high, the application rates of all herbicides were increased accordingly. Any extrapolation of this work will therefore need to consider these factors, plus rainfall in deciding on the herbicide rates which will be effective, economic and safe.

ACKNOWLEDGEMENTS

Appreciation is expressed to growers who provided the trial sites, and to Ruakura Soil Testing Laboratory for the soil analyses.

REFERENCES

- Dawbin, N.T., 1980. Asparagus — cultural recommendations. *N.Z. Cropping News*, January, 1980.
- Franklin, S.J., 1980 Weed control in asparagus. *Proc. seminar on asparagus, Hamilton*: 37-41, Advisory Services Division, MAF, Hamilton.