Horticultural Crops

DAMAGE TO PASSIONFRUIT AND STRAWBERRY CAUSED BY THE AUSTRALIAN CROP MIRID

R. T. BAKER

*Horticultural Research Centre, Research Division,
Ministry of Agriculture and Fisheries, Levin.*

Summary

In 1976 damage to young fruit and shoots of passionfruit (*Passiflora* sp) in the Levin district was caused by adults of the Australian crop mirid (*Sidnia kinbergi*). Laboratory trials showed that this mirid is readily controlled by maldison and dichlorvos but under field conditions there is rapid reinfestation from surrounding pasture. During 1978 nymphs and adults of the Australian crop mirid were found to damage flowers and fruit of strawberry (*Fragaria x ananassa*). A heavy infestation was checked by an application of mevinphos but reinfestation from surrounding clover occurred within a fortnight.

INTRODUCTION

The Australian crop mirid (*Sidnia kinbergi* [Stal, 1859]) has been more widely known in the literature as *Eurystylus australis* Poppius, 1911, and all New Zealand references to this insect are under the junior synonym.

Myers (1926) mentioned very serious damage to passion vine shoots and also noted that the insect had been taken on apple and frequently on grasses. Cumber (1959) found small numbers of nymphs and adults during a survey of pastures in the North Island but concluded that, although widespread, the mirid was not of economic significance. Eyles (1960) recorded the Australian crop mirid more frequently on fodder crops than on pasture and mentioned its potential as a virus or fungus vector through its feeding habits. A recent reference by Ferro (1976) mentions the Australian crop mirid as the dominant insect in lucerne stands but the extent of damage is unknown.

This paper confirms the damage to passionfruit, records for the first time serious damage to strawberry fruit and flowers, and notes the abundance of the Australian crop mirid on clover swards.

DAMAGE

In February 1976 a commercial crop of passionfruit near Levin was examined after the grower noticed damage on skin of immature fruits. Damage was initially in the form of 'stings' which formed disfiguring callouses as the fruit matured. As many as 26 stings were found on a single young fruit. Net sweepings of the passion vines revealed a large number of mirids. The mirids were also taken in abundance from the surrounding headlands and especially from the clover sward between the rows of vines. In the insectary adult mirids were caged in small tubes, stoppered with young passionfruits or shoots. Some were also kept in large cages with access to flowers, fruits and shoots. The mirids soon pierced the skin and fed on the young fruits or stems, causing the damage seen in the field.

Proc. 31st N.Z. Weed and Pest Control Conf.

Horticultural Crops

Badly affected young shoots withered and dropped from the plant but in the field damage was confined to fruit. Damaged fruit continued to grow and mature with only the appearance being affected.

Further net sweepings of passion vines were made in January and February 1978 when many adult mirids were collected. No nymphs have been found on passionfruit, even after very thorough hand searching.

Deformed strawberry fruits and blackened flowers were first discovered at HRC, Levin, on 31.1.78. Hand and net searches showed very large numbers of nymphs and adults of the Australian crop mirid on strawberry and surrounding clover. Insectary observations confirmed that damage observed in the field was caused by this insect. Flowers which had turned black did not produce fruit while affected fruits were badly distorted and unmarketable. In the laboratory adults and nymphs seemed to prefer to feed on strawberry fruits but they also fed on strawberry flowers and stems as well as on clover seed heads, flowers and stems.

DISTRIBUTION AND LIFE HISTORY

The Australian crop mirid occurs widely throughout the North Island and the north of the South Island (Eyles 1960) and is extremely common in the Horowhenua district. Observations in Levin confirm the reports of Myers (1926), Cumber (1959) and Eyles (1960) that adults are present from January to April but no details of the life history are known. Most Miridae insert their eggs into the softer parts of plants (Miller 1956) but so far microscopic examination of clover and strawberry shoots has failed to reveal eggs of the Australian crop mirid. From wing and body length measurements of field collected samples it appears that there are five nymphal stages. In the laboratory nymphal stages lasted for only 2-3 days while adults lived for about 7 days.

CONTROL

In an insectary trial fully open passionfruit flowers were placed in water and enclosed in a plastic plant propagator. The flowers were dipped in maldison (2 g/litre) or dichlorvos (2.27 g/litre) or were untreated. After the flowers had been dipped and spray deposit allowed to dry four adult mirids were introduced into each propagator. There were six replications of each treatment. After 24 hours all mirids in the maldison and dichlorvos cages were dead but there was no mortality in the untreated controls.

On passionfruit, growers have used parathion, TDE and azinphos-methyl, with apparently little effect, while mevinphos applied to strawberry cleared the area of mirids for about 14 days. The data collected from the insectary trial and the field applications show that organophosphates will readily kill the mirids but reinfestation is rapid and it would be necessary to spray surrounding clover swards to have any lasting benefit. Possibly control measures could be improved if the mirid's life cycle was known in detail. Spraying with organophosphates at full flower should be avoided because of possible damage to bee populations.

CONCLUSION

The Australian crop mirid is probably of little economic significance to the passionfruit grower as damage is to the skin of the fruit only and does not seem to diminish the crop. On strawberry, however, the mirid poses a distinct threat and considerable losses could occur. Organophosphate insecticides give control but reinfestation, particularly from clover, must be watched for.

Horticultural Crops

ACKNOWLEDGEMENTS

The identification by Prof. T. E. Woodward, University of Queensland, the co-operation of the grower Mr J. Wylie of Muhunua East, and the assistance with field work of Mrs J. Batten, Levin HRC, are acknowledged with thanks.

REFERENCES

- Cumber, R. A., 1959. The insect complex of sown pastures in the North Island. II. The Hemiptera as revealed by summer sweep-sampling. *N.Z.J. Agric. Res.* 2: 1-25.
- Eyles, A. C., 1960. Insects associated with the major fodder crops in the North Island. II. Hemiptera. *N.Z.J. Agric. Res.* 3: 994-1008.
- Ferro, D. N., 1976. In New Zealand insect pests, ed. D. N. Ferro, Lincoln University, College of Agriculture, Canterbury, N.Z., 130.
- Miller, N. C. E., 1956. The biology of the heteroptera. Leonard Hill Books Ltd, London.
- Myers, J. G., 1926. Biological notes on the New Zealand heteroptera. *Trans. N.Z. Inst.*: 56: 449-511.