SPEAR SPOT OF ASPARAGUS

L.H. CHEAH

Horticultural Research Centre, MAF, Levin

Spear spot of asparagus, caused by *Fusarium redolens*, was first recorded in New Zealand in 1965 (Dingley 1965). Since then there have been no reports of the disease, probably because it has not caused significant crop damage. However, in the last two seasons the disease has become more evident and on one property, caused total rejection of a consignment of asparagus. The disease was not controlled by the routine spray programme for use on asparagus. This note describes spear spot of asparagus and outlines studies on the pathogenicity and fungicide control of *F. redolens*.

Pathogenicity

Spear spot appears as minute, brown to rusty brown spots at the tips of bracts on young asparagus spears. Under humid conditions, most of the bracts on a spear become infected and traces of white fungal mycelium can be observed on these spots. Spots found on spear stems are larger (2-5 mm) and usually occur at ground level. In severe cases, lesions occur further up causing the spears to be deformed and stunted.

Fungi were isolated from field grown diseased asparagus spears onto agar media and after subculturing, they were identified as *F. redolens*. Pathogenicity was tested using asparagus spears, injured by pricking with a sharp scalpel on stem and bracts and inoculated with spore and mycelial suspensions of the fungus. The spears were then placed in a sealed plastic box to maintain high humidity and held at room temperature ($21 \pm 3^\circ\text{C}$). Typical symptoms of rusty brown spots seen in the field were observed on bracts and stems of spears 5 days after inoculation. Uninoculated spears remained disease-free. *F. redolens* was isolated from inoculated spears.

An investigation into effects of weather conditions on disease development was carried out. Percentage of infected spears (200 spears/sample) was recorded in 10 harvests during October 1985, and results were compared with meteorological records (rainfall, soil and air temperatures, and humidity) taken at Levin Horticultural Research Centre, 4 km from the harvest site. Between 0 and 16% of spears showed spear spot symptoms at respective harvests but no obvious relationship between disease incidence and meteorological records could be established.

TABLE 1: Effect of fungicides on growth of *Fusarium redolens* on culture plates and asparagus spears.

Fungicide		Poison agar plate (colony diam. mm)	Spear technique (lesion length mm)	
Common name (ppm)	Trade name		Surface application	Systemic soak
1 benomyl 500	Benlate 50WP	0*	17.0	0
2 captan 1000	Othocide 80WP	23.2	29.4	30.0
3 iprodione 500	Rovral 50WP	30.2	30.2	26.2
4 mancozeb 1500	Manzate 80WP	16.4	27.8	30.8
5 metalaxyl MZ 1500	Ridomil MZ72	21.4	30.4	27.6
6 prochloraz 500	Sportak 50WP	0	22.6	19.2
7 thiram 1500	Thiram 80WP	18.2	28.0	29.6
8 untreated control		64.2	30.4	32.0
LSD (P = 5%)		1.4	3.6	4.7

*Zero result was not included in the analysis of variance.

Proc. 39th N.Z. Weed and Pest Control Conf.

Fungicide Screening

Two techniques were used to test fungicides in the laboratory:

- (a) Agar plate technique. Fungicides (Table 1) were added to cooled molten potato dextrose agar (PDA) to give required concentrations (Pennycook and Corbin 1970) and agar plates were prepared. An inoculum plug (6 mm diam.) was placed, mycelium surface downwards, in the centre of each plate, and the plates were incubated at 20 °C. Five plates were used for each fungicide. After 6 days, the diameters of resulting colonies were measured.

Benomyl and prochloraz completely inhibited mycelial growth (Table 1). The other fungicides were less effective, but all reduced growth of *F. redolens*.

- (b) Asparagus spear technique. Healthy field-grown asparagus spears (2-3 days old) were washed in tap water and used for screening fungicides by the following techniques;

- (i) Surface application: Spears were sprayed with fungicides (Table 1) to run off, left to air-dry and trimmed to 10 cm long. Thin slivers of epidermal tissue were removed along their lengths before placing them in plastic containers. An inoculum plug (6 mm) from a 5-day-old PDA culture was placed, mycelium surface downwards, onto the centre of the wound of each spear. Five spears were used for each fungicide treatment. The containers were then sealed to maintain high humidity and incubated at 20 °C. Lengths of disease lesions were measured after 6 days. A similar procedure was used for controls with spears sprayed with water only.
- (ii) Systemic treatment: Five spears were immersed to about 2 cm in a beaker containing fungicide suspension. Separate beakers were used for each fungicide (Table 1), and were continuously shaken to maintain the fungicides in suspension. Spears were removed 4 h later and dried with a paper towel. The upper portions were then cut to about 10 cm length, and injured, inoculated, incubated and recorded as described above. Control spears were soaked in water instead of fungicides.

Benomyl completely prevented disease development with the systemic soak technique (Table 1). Prochloraz was less effective but still provided good disease control. With surface application, benomyl was again better than prochloraz. Of the other fungicides, only iprodione slightly inhibited lesion development ($P < 0.05$).

Spear spot of asparagus caused by *F. redolens* is a disease of potential economic importance. Benomyl and prochloraz were shown to be very effective for disease control in laboratory tests, and their use should be further evaluated in the field. Asparagus spear techniques have proved to be rapid and reliable for pathogenicity testing and fungicide screening in the laboratory. They also provide a better simulation of the field situation than do agar methods and may therefore give more practical results.

ACKNOWLEDGEMENTS

I thank Drs E.H.C. McKenzie and W.F.T. Hartill of Plant Diseases Division, DSIR, Auckland for fungal identification and technical advice respectively. Field assistance was provided by N. Mark.

REFERENCES

- Dingley, J.M., 1965. New records of fungus disease of plants in New Zealand, 1962-64. *N.Z. J. Agric. Res.* 8: 905-920.
- Pennycook, S.R. and Corbin, J.B., 1970. A preliminary laboratory assessment of fungicide for the control of *Sclerotinia sclerotiorum* (Lib.) al By. *Tane* 16: 21-30.