RESISTANCE TO BENOMYL IN ISOLATES OF THE EYESPOT FUNGUS IN CEREALS IN SOUTHLAND

A.C. KING¹, M.G. CROMEY², F.R. SANDERSON³
and P.R. SCOTT⁴

¹ Botany Department, University of Canterbury, Christchurch

² Plant Diseases Division, DSIR, Lincoln

³ Crop Research Division, DSIR, Lincoln

⁴ Plant Breeding Institute, Trumpington, Cambridge, England

SUMMARY

A survey in Southland during February 1984 has found resistance to benomyl present in many isolates of the cereal eyespot pathogen *Pseudocercospora herpotrichoides*. Similar results have been found in surveys in Europe. Isolates were found to vary in resistance to benomyl and were present in both barley and wheat crops, these often having histories of benomyl usage.

INTRODUCTION

Pseudocercospora herpotrichoides (Fron) Deighton, the causal agent of the cereal eyespot disease, is a fungus which infects the base of the culms of the host plants. Characteristic symptoms include the formation of ellipsoid lesions with darkened edges and inner spots resembling eyes. Severe infection leads to stem lodging, shrivelled grain and partially empty ears. The eyespot mycelium persists between crops on cereal debris in the soil. Control may be by crop rotation, cultural methods and by suitable fungicide usage (Harvey and Close 1981).

Benzimidazole fungicides have been used successfully over the past decade in many areas of the world. However, isolates of *P. herpotrichoides* with resistance to benomyl and related benzimidazole fungicides have been reported in Europe, including some from paddocks never treated with these fungicides (Rashid and Schlosser 1977; Horsten and Fehrmann 1980; Brown *et al* 1984); this has raised doubts regarding the continued effectiveness of benzimidazole fungicides in controlling eyespot. An unpublished survey of isolates from infected cereal crops in the UK in 1982 and 1983 showed that more than 40 per cent were resistant to 2 µg/ml of benomyl in culture (M.J. Griffin, pers comm).

Two distinct types of *P. herpotrichoides* have been characterised by European studies, these being termed R-type and W-type (Lange de la Camp 1966; Scott *et al* 1975). R-type isolates are equally pathogenic to wheat and rye and on potato dextrose agar (PDA) have a slow growth rate and a 'feathery' colony edge. W-type isolates are more pathogenic to wheat than rye, have a faster growth rate and an 'even' colony edge on PDA. UK results suggest that benomyl-resistant isolates are more frequently R-type than W-type (M.J. Griffin pers comm; Brown *et al* 1984).

In New Zealand, eyespot is chiefly important on spring-sown wheat and barley in Southland, where high rainfall, frequent cereal cropping and susceptible varieties promote infection, even though the disease is more typically a problem on autumn-sown cereals. The value of benomyl for controlling eyespot in Southland was recognised in the early 1970's (Witchalls and Close 1971) and extensive use of the fungicide for this purpose started somewhat earlier than in the UK.

It was considered important to survey isolates of eyespot from Southland for resistance to benomyl (Benlate 50W). This paper reports the results of a survey made during February 1984. Sampled crops were in most cases late-maturing. All crops were between growth stages 71 and 79 (Zadocks *et al* 1974) when sampled. Some tests were also made for resistance to prochloraz, an alternative fungicide known to control eyespot.

Proc. 37th N.Z. Weed and Pest Control Conf.

MATERIALS AND METHODS

Survey

A wide range of cereal paddocks was sampled. Where possible paddock histories and previous fungicide applications were determined. In each paddock, 10 random samples were made of at least 10 tillers, each being reasonably spaced. If a crop was partially lodged then both lodged and unlodged tillers were sampled. Tillers were uprooted, headed and the leaf sheaths removed.

Isolation

Stems were surface sterilised in 1-2 per cent sodium hypochlorite for 2 minutes. Sections were sliced from lesions on the stem or pieces of mycelium were taken from the inner stem lumens. These were plated onto antibiotic-amended PDA and incubated at 18 °C. Subcultures were made onto non-amended PDA within 2-3 days to rescue isolates from faster growing fungi such as *Fusarium* spp.

Fungicide-sensitivity tests

Isolates were tested for resistance by plating onto PDA amended with 2 µg/ml benomyl. Ten selected isolates were later tested over a range of benomyl concentrations. Prochloraz (Sportak 45EC) sensitivity was tested at 2 µg/ml for 22 isolates.

RESULTS AND DISCUSSION

Of 53 paddocks sampled, *P. herpotrichoides* was isolated from 43. The one rye paddock provided two benomyl-sensitive isolates. Of the 19 barley paddocks, five contained resistant isolates and of the 23 wheat paddocks, seven contained resistant isolates. Table 1 gives the percentage of benomyl-resistant isolates in a total of 131 isolates.

Table 2 presents the paddocks within which benomyl-resistant isolates were found and gives the proportion of such isolates.

TABLE 1: Sensitivity to 2 µg/ml benomyl of *P. herpotrichoides* isolated from rye, wheat and barley in Southland in 1984.

Isolate	Isolated from			Total
	Rye	Wheat	Barley	
Sensitive	2 (100%)	45 (81.81%)	62 (83.78%)	109
Insensitive		10 (18.18%)	12 (16.22%)	22

TABLE 2: Occurrence of benomyl resistance in 12 cereal crops in Southland in 1984.

Paddock	Location	Crop	Previous crop	Benomyl sprays	Total Number of isolates	No. of Resistant isolates
7b	Otahuti	Barley	2 wheat	3	5	5
27	Gridirons	Wheat	1 wheat	2	1	1
35	South Kelso	Wheat			1	1
41	Near Conical Hill	Wheat			2	2
45	Waitane	Wheat			7	1
46	Waitane	Wheat			4	1
47	Waitane	Wheat			6	2
50	Hedgehope	Barley	2 wheats	2	5	3
52	Hedgehope	Barley	2 wheats	2	3	1
53	Hedgehope	Barley	Oats: 5 wheat	5	1	1
59	Winton	Wheat	5 wheat	5	3	2
60	Winton	Barley			6	2

Fig. 1: Percentage of isolates of *P. herpotrichoides* growing at various benomyl concentrations.

Paddocks 7a and 7b were neighbouring sites at Otahuti. Both were first-year barley crops which had been sprayed with half-rates of benomyl. Paddock 7a had a history of 14 years of non-cereal crops and contained insignificant eyespot infection, while paddock 7b had been preceded by two benomyl-sprayed wheat crops following 12 years out of cereals. The barley crop in paddock 7b was severely infected with eyespot and all isolates were benomyl resistant. As paddock 7a was a first-year cereal it was expected that eyespot infection would be minimal. Paddock 7b exemplifies how rapidly eyespot inoculum develops and how benomyl resistance probably occurs as a result of selection pressures.

Paddocks 53 and 59 had crop histories of at least four years of cereals and benomyl applications and contained benomyl-resistant eyespot isolates.

It is unknown if New Zealand paddocks which have been untreated with benomyl fungicides have benomyl-resistant isolates within the wild populations.

The 10 isolates examined in detail were found to differ in the maximum concentration of benomyl tolerated (Fig. 1). All grew normally at concentrations up to 0.2 µg/ml. Of those that resisted 2 µg/ml, many also resisted 20 and 200 µg/ml. Brown *et al* (1984) reported that many UK isolates were sensitive to the related compound carbendazim even at concentrations below 0.1 µg/ml, while some resistant isolates grew at 1000 µg/ml.

All the 22 isolates tested (including benomyl-sensitive and benomyl-resistant isolates) were sensitive to 2 µg/ml of prochloraz.

Isolates conforming morphologically to both W-type and R-type forms were isolated in Southland, but intermediate forms also appeared to be present, making precise classification difficult. Relative pathogenicity to rye of the different New Zealand forms is not known. In Southland, most benomyl insensitive isolates were fast growing in contrast to the UK, where most are slow growing (R-type) isolates (M.J. Griffin, pers comm).

The presence of benomyl resistance in Southland is a cause for concern, although the impact on eyespot control has yet to be determined. In the UK, the occurrence of resistant isolates has been positively associated with ineffective eyespot control by

benzimidazole fungicides in crops, although their use may still give cost-effective yield improvement (M.J. Griffin and D.T. Yarham pers comm). Changes in the frequency of resistance in the pathogen population must be monitored and the effect of this on control of eyespot with benzimidazole fungicides assessed.

REFERENCES

- Brown, M.C., Taylor, G.S. and Epton, H.A.S., 1984. Carbendazim resistance in the eyespot pathogen. *Pseudocercospora herpotrichoides*. *Plant Pathology* 33: 101-111.
- Harvey, I.C. and Close, R.C., 1981. Eyespot of Wheat. Symptoms and Control. Aglink FPP 236 (2nd revise).
- Horsten, J. and Fehrmann, H., 1980. Fungicide resistance of *Septoria nodorum* and *Pseudocercospora herpotrichoides*. 1. Effect of fungicide application on frequency of resistant spores in the field. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz*. 87: 439-453.
- Lange-de la Camp M., 1966. Die Wirkungsweise von *Cercospora herpotrichoides* Fron, der Erreger der Halmbruchkrankheit des Getreides. II. Aggressivität des Erregers. *Phytopathologische Zeitschrift* 56: 155-190.
- Rashid, T. and Schlosser, E., 1977. Cereal foot rot pathogens. II. Occurrence and distribution of benomyl-tolerant strains. *Mededelingen van de Faculteit der Landbouwwetenschappen Rijksuniversiteit Gent* 42: 1057-1065.
- Scott, P.R., Hollings, T.W. and Muir, P., 1975. Pathogenicity of *Cercospora herpotrichoides* to wheat, barley, oats and rye. *Transactions of the British Mycological Society* 65: 529-538.
- Witchalls, J.T. and Close, R., 1971. Control of eyespot lodging in wheat by benomyl. *Plant Disease Reporter* 55: 45-47.
- Zadoks, J.C., Chang, T.T. and Konzac, C.F., 1974. A decimal code for the growth stages of cereals. *Weed Research* 14: 415-421.