THE EFFECT OF GREENHOUSE WHITEFLY SPRAY THRESHOLDS ON TAMARILLO YIELD

R.H. BLANK

Agricultural Research Division, MAF, P.O. Box 943, Whangarei

SUMMARY

Tamarillo (*Cyphomandra betacea*) infested with greenhouse whitefly (*Trialeurodes vaporariorum*) were sprayed with an oil-pyrethroid mixture at three threshold levels determined by adult densities on young leaves. Whitefly on unsprayed tamarillos caused considerable leaf and fruit drop and severe contamination with sooty mould. Fruit was smaller and fruit yield reduced by 43%. The 20 adults per leaf threshold required 12 spray applications at an average interval of 8 days which prevented sooty mould contamination of leaves and fruit. The 50 and 100 adults per leaf threshold treatments suffered sooty mould contamination, 17-39% defoliation, and 19-30% reduction in yield compared with the lower threshold treatment.

INTRODUCTION

The greenhouse whitefly is a serious pest of tamarillo. Adult whitefly congregate on the underside of young leaves on which they feed and lay eggs. The first instar stages hatch on the young leaves, with older larval and pupal stages progressively found on older, mature leaves. The whitefly contaminate leaves and fruit with a honeydew secretion on which grows a black sooty mould. In severe infestations, fruit and leaf drop occurs (Blank *et al* 1986a).

Pyrethroid sprays, either alone or in combination with oil, can give high mortalities of all stages (Blank *et al* 1986b). Control of whitefly within a tamarillo orchard is difficult to maintain because of the difficulty in achieving complete spray coverage of the dense foliage and reinvasion by adults. The frequency of spraying required for control depends on the rate of development of whitefly which varies with temperature. An adult threshold count based on the selection of young leaves at the top of the canopy was found to be a useful technique for determining when to spray (Blank *et al* 1986a).

This study investigates suitable spray thresholds to prevent crop loss.

MATERIALS AND METHODS

An 0.5 ha block of 5 year old red tamarillos in Kerikeri was used for this trial. Plant spacing was 1 m with 4.5 m between rows. Each plot comprised eight tamarillo plants. All assessments were made from the four centre plants. Buffer rows were alternated with rows containing treatment plots and were sprayed only on 14 March and 17 April 1986 when invasion of whitefly onto sprayed plots became difficult to control.

All insecticide applications used a mixture of deltamethrin (Decis 2.5 EC 0.6 g ai/100 litres) and oil (Sunspray mineral oil 96 EC 300 g ai/100 litres). The whole tamarillo block was sprayed, using an airblast sprayer, 3 weeks prior to the first threshold count on 30 January 1986 to bring whitefly populations to a uniform level.

The trial site was visited 15 times at 5-12 day intervals until 8 May. Adult counts were taken from each plot and a decision to spray individual plots made. Threshold counts were made from five leaf tips (70-150 mm long) at the top of the canopy which were quickly selected as having high adult densities (Blank *et al* 1986). Individual plots were sprayed to the point of run off with a motorised knapsack sprayer (setting 3).

Four treatments (untreated, 20, 50 and 100 adults/leaf threshold) were assigned at random amongst 24 plots. Some plots markedly exceeded the assigned threshold treatment between visits. Where this occurred, they were re-allocated to a higher threshold treatment on 14 March so that at the completion of the trial there were four

Proc. 40th N.Z. Weed and Pest Control Conf.

plots with a 20 adults/leaf threshold treatment, seven plots with 50 and 100 adults/leaf threshold treatments and six untreated plots.

Sooty mould was assessed for each plot at each visit. Fruit and leaves were ranked on a scale of 0 (nil) to 5 (very heavy contamination). Leaf length and new leaves produced in front of 10 tagged young leaves on unsprayed plants were recorded at each visit from 14 March until 1 May. Defoliation was assessed on 7 May from eight shoots per plot. Leaf abscission scars from the third leaf below the growing tip to the 23rd leaf site were counted. Live larvae and pupae were assessed from 40 leaves picked from each plot on 28 May. Ripe fruit were picked on 7 and 18 May and all remaining fruit strip-picked on 3 June. Fruit on the ground under assessment trees were also collected at these times. Fruit were bulk weighed and a sub-sample of 10 fruit/plot individually weighed and assessed for sooty mould.

RESULTS

Adult whitefly on the unsprayed tamarillo plots underwent an exponential increase up to late February followed by a slower rate of increase until early May when densities decreased (Fig. 1). The 20 adults/leaf threshold treatment required 12 spray applications at an average interval of 8 days and contained whitefly to a cumulative threshold count of 47 (Table 1, Fig. 1). The 50 and 100 adults/leaf threshold treatments received fewer spray applications and cumulative threshold counts were intermediate between the unsprayed and the 20 threshold treatment. Larval and pupal densities in May followed similar trends to the cumulative adult threshold counts (Table 2).

Fig. 1: Cumulative adult whitefly counts at each visit for unsprayed, 100, 50 and 20 adults per leaf spray thresholds.

TABLE 1: The effect of adult whitefly spray thresholds on whitefly infestations, spray intervals, defoliation, sooty mould and fruit yield on tamarillo.

	Whitefly threshold (adults/leaf)				Av. SEM
	20	50	100	Unspr.	
Cumulative threshold (adults/leaf)	47	98	125	217	16
No. sprays	12	9	8	—	—
Spray intervals (days)	8	10	12	—	—
Defoliation (%)	15	32	54	93	6
Sooty mould — leaves	nil	moderate	heavy	v. heavy	—
fruit	nil	light	light	v. heavy	—
Fruit size (g)	71	70	71	61	2
Fruit drop (kg/m of row)	0.5	0.4	0.4	1.2	0.2
Fruit yield (kg/m of row)	8.0	5.6	6.5	3.4	0.6

TABLE 2:: The effect of adult whitefly spray thresholds on percentage of tamarillo leaves infested with larvae and pupae.

Larvae and pupae/leaf =	% leaves infested Whitefly threshold (adults/leaf)			Unspr.
	20	50	100	
0	93	69	60	0
1-20	5	19	16	0
21-100	2	11	7	0
101-1000	0	1	7	7
1001-20000	0	0	0	93

Fig. 2: Tamarillo leaf growth measured as leaf length and new leaves produced.

Rapid tamarillo growth, as measured by leaf extension and the production of new leaves, occurred in March (Fig. 2). Leaf production slowed in response to high whitefly densities and/or drought conditions in April and May.

The high densities of whitefly on the unsprayed tamarillos caused severe defoliation of plants, contaminated both leaves and fruit with heavy deposits of sooty mould, caused considerable fruit drop, reduced fruit size and reduced fruit yield to 43% of that in the 20 adult/leaf threshold treatment (Table 1). Sooty mould developed rapidly on leaves and fruit of unsprayed plants with contamination light in late January, heavy by late February and very heavy by mid-March. The 50 and 100 threshold treatments also suffered leaf defoliation and sooty mould contamination and yields were reduced to 70 and 81% respectively of that in the 20 threshold treatment (Table 1). A very light sooty mould contamination of leaves had occurred on half of the 50 and 100 threshold plots by late January. By late February there was contamination of leaves and fruit from half of the plots. Moderate to heavy contamination of leaves occurred after late March with fruit contamination remaining light and not all plots affected. Significant regression relationships were found between fruit yield and cumulative whitefly threshold counts, number of sprays and percent defoliation (Table 3).

TABLE 3: The regression relationship of tamarillo fruit yield (g/m of tamarillo row) versus whitefly threshold (adults/leaf), number of sprays and percent defoliation.

	Slope	Intercept	r ²	Significance
Cumulative threshold	-22.5	8.6	0.45	P<0.001
No. of sprays	362	3.1	0.66	P<0.001
% defoliation	-41	7.8	0.39	P<0.01

DISCUSSION

Regularly sprayed tamarillo treatments were invaded by adult whitefly from neighbouring unsprayed tamarillos. Adults rapidly reinfested the vigorous new growth on the regularly sprayed tamarillos which was produced perhaps even faster than the one leaf every 8.5 days that occurred in March on the unsprayed tamarillos. This explains why spray intervals need to be close during periods of rapid growth.

The rapid whitefly reinfestation combined with the 5 or more days or greater interval between visits resulted in the 20 adults/leaf threshold treatment having a cumulative threshold close to 50 adults/leaf. This overshooting of the threshold level also occurred in a trial carried out in adjacent commercial blocks sprayed at a threshold of 50 adults/leaf (Blank *et al* 1986a). The spray interval on the commercial blocks of 10 days was similar to the 9 days found for the 50 adults/leaf threshold treatment in this trial, although both would have been influenced by assessment interval. Ideally, whitefly populations should be assessed about 5 days after insecticide applications in warm weather and then at intervals dependent on the previous threshold count.

Whitefly infestations, if left unchecked, can have a catastrophic effect on defoliation, fruit size and yield as well as causing heavy sooty mould contamination. Treatments with spray thresholds of 50 and 100 adults, although containing populations at relatively low levels, still suffered considerable defoliation and yield losses. In addition, the light sooty mould contamination would still necessitate washing and down-grading of fruit.

The commercial viability of tamarillo blocks is dependent on achieving good yields. Whitefly infestations must be contained at low levels to prevent crop loss. A threshold count of approximately 20 appears necessary to prevent both sooty mould contamination of fruit and yield loss.

ACKNOWLEDGEMENTS

My thanks to Miss M.J. Blagrove, Messrs M.H. Olson and D.S. Bell for technical assistance and to Mr A.R.W. White for use of the tamarillo block.

REFERENCES

- Blank, R.H., Blagrove, M.J. and White, A.R.W., 1986a. Strategies for control of greenhouse whitefly in tamarillo orchards. *Proc. 39th N.Z. Weed and Pest Control Conf.*: 153-157.
- Blank, R.H., Blagrove, M.J. and Waller, J.E., 1986b. Screening chemicals to control greenhouse whitefly on tamarillo. *Proc. 39th N.Z. Weed and Pest Control Conf.*: 148-152.